

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/12, C07K 14/47, C12N 15/62, C07K 16/18, C12N 5/10</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/63377</b> (43) International Publication Date: <b>26 October 2000 (26.10.00)</b>
(21) International Application Number: <b>PCT/US00/10454</b> (22) International Filing Date: <b>19 April 2000 (19.04.00)</b> (30) Priority Data: <b>09/294,943                      20 April 1999 (20.04.99)                      US</b> (71) Applicant: <b>ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US).</b> (72) Inventors: <b>PIDDINGTON, Christopher, S.; 2838 Irongate Place, Thousand Oaks, CA 91362 (US). BISHOP, Paul, D.; 28425 S.E. 8th Street, Fall City, WA 98024 (US).</b> (74) Agent: <b>LINGENFELTER, Susan, E.; ZymoGenetics, Inc., 1201 Eastlake Avenue East, Seattle, WA 98102 (US).</b>			(81) Designated States: <b>AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: <b>ADIPOCYTE COMPLEMENT RELATED PROTEIN HOMOLOG ZACRP3</b>			
(57) Abstract  The present invention relates to polynucleotide and polypeptide molecules for zacrp3, a novel member of the family of proteins bearing a collagen-like domain and a C1q domain. The polypeptides and polynucleotides encoding them, are involved in dimerization or oligomerization and may be used in the study thereof. The present invention also includes antibodies to the zacrp3 polypeptides.			

**A10 - 09/997,610**

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5       ADIPOCYTE COMPLEMENT RELATED PROTEIN HOMOLOG ZACRP3

## BACKGROUND OF THE INVENTION

Energy balance (involving energy metabolism, nutritional state, lipid storage and the like) is an  
10 important criteria for health. This energy homeostasis involves food intake and metabolism of carbohydrates and lipids to generate energy necessary for voluntary and involuntary functions.. Metabolism of proteins can lead to energy generation, but preferably leads to muscle  
15 formation or repair. Among other consequences, a lack of energy homeostasis lead to over or under formation of adipose tissue.

Formation and storage of fat is insulin-modulated. For example, insulin stimulates the transport  
20 of glucose into cells, where it is metabolized into  $\alpha$ -glycerophosphate which is used in the esterification of fatty acids to permit storage thereof as triglycerides. In addition, adipocytes (fat cells) express a specific transport protein that enhances the transfer of free fatty  
25 acids into adipocytes.

Adipocytes also secrete several proteins believed to modulate homeostatic control of glucose and lipid metabolism. These additional adipocyte-secreted proteins include adipsin, complement factors C3 and B,  
30 tumor necrosis factor  $\alpha$ , the ob gene product and Acrp30. Evidence also exists suggesting the existence of an insulin-regulated secretory pathway in adipocytes. Scherer et al., J. Biol. Chem. 270(45): 26746-9, 1995. Over or under secretion of these moieties, impacted in  
35 part by over or under formation of adipose tissue, can lead to pathological conditions associated directly or indirectly with obesity or anorexia.

Acrp30 is a 247 amino acid polypeptide that is expressed exclusively by adipocytes. The Acrp30 polypeptide is composed of a amino-terminal signal sequence, a 27 amino acid stretch of no known homology, 22 perfect Gly-Xaa-Pro or imperfect Gly-Xaa-Xaa collagen repeats and a carboxy terminal globular domain. See, Scherer et al. as described above and International Patent Application No. WO 96/39429. Acrp30, an abundant human serum protein regulated by insulin, shares structural similarity, particularly in the carboxy-terminal globular domain, to complement factor Clq and to a summer serum protein of hibernating Siberian chipmunks (Hib27). Expression of Acrp30 is induced over 100-fold during adipocyte differentiation. Acrp30 is suggested for use in modulating energy balance and in identifying adipocytes in test samples.

Another secreted protein that appears to be exclusively produced in adipocytes is apM1, described, for example, in Maeda et al., Biochem. Biophys. Res. Comm. 221: 286-9, 1996. A 4517 bp clone had a 244 amino acid open reading frame and a long 3' untranslated region. The protein included a signal sequence, an amino-terminal non-collagenous sequence, 22 collagen repeats (Gly-XAA-Pro or Gly-Xaa-Xaa), and a carboxy-terminal region with homology to collagen X, collagen VIII and complement protein Clq.

Complement factor Clq consists of six copies of three related polypeptides (A, B and C chains), with each polypeptide being about 225 amino acids long with a near amino-terminal collagen domain and a carboxy-terminal globular region. Six triple helical regions are formed by the collagen domains of the six A, six B and six C chains, forming a central region and six stalks. A globular head portion is formed by association of the globular carboxy terminal domain of an A, a B and a C chain. Clq is therefore composed of six globular heads linked via six collagen-like stalks to a central fibril region. Sellar et al., Biochem. J. 274: 481-90, 1991. This configuration

is often referred to as a bouquet of flowers. Acrp30 has a similar bouquet structure formed from a single type of polypeptide chain.

Clq has been found to stimulate defense mechanisms as well as trigger the generation of toxic oxygen species that can cause tissue damage (Tenner, Behring Inst. Mitt. 93:241-53, 1993). Clq binding sites are found on platelets. Additionally complement and Clq play a role in inflammation. The complement activation is initiated by binding of Clq to immunoglobulins

Inhibitors of Clq and the complement pathway would be useful for anti-inflammatory applications, inhibition of complement activation and thrombotic activity.

The present invention provides such polypeptides for these and other uses that should be apparent to those skilled in the art from the teachings herein.

#### SUMMARY OF THE INVENTION

Within one aspect, the invention provides an isolated polypeptide comprising a sequence of amino acid residues that is at least 75% identical in amino acid sequence to residues 51-246 of SEQ ID NO:2, wherein the sequence comprises: Gly-Xaa-Xaa or Gly-Xaa-Pro repeats forming a collagen domain, wherein Xaa is any amino acid; and a carboxyl-terminal Clq domain comprising 10 beta strands. Within one embodiment the polypeptide that is at least 90% identical in amino acid sequence to residues 23-246 of SEQ ID NO:2. Within another embodiment the collagen domain consists of 15 Gly-Xaa-Xaa repeats and 6 Gly-Xaa-Pro repeats. Within another embodiment the carboxyl-terminal Clq domain comprises the sequence of SEQ ID NO:10. Within another embodiment the carboxy-terminal Clq domain comprises amino acid residues 119-123, 140-142, 148-151, 155-158, 161-173, 175-182, 190-197, 200-212, 217-222 and 236-241 of SEQ ID NO:2. Within another embodiment

any differences between the polypeptide and SEQ ID NO:2 are due to conservative amino acid substitutions. Within another embodiment the polypeptide specifically binds with an antibody that specifically binds with a polypeptide consisting of the amino acid sequence of SEQ ID NO:2. Within still another embodiment the polypeptide comprises residues 23-246 of SEQ ID NO:2. Within another embodiment the collagen domain consists of amino acid residues 51-113 of SEQ ID NO:2. Within yet another embodiment the C1q domain consists of amino acid residues 114-246 of SEQ ID NO:2. Within another embodiment the polypeptide is covalently linked at the amino or carboxyl terminus to a moiety selected from the group consisting of affinity tags, toxins, radionucleotides, enzymes and fluorophores.

Also provided is an isolated polypeptide selected from the group consisting of: a) a polypeptide consisting of a sequence of amino acid residues that is 80% identical in amino acid sequence to amino acid residue 51 to amino acid residue 113 of SEQ ID NO:2, the polypeptide consisting of Gly-Xaa-Xaa and Gly-Xaa-Pro repeats forming a collagen domain; b) a polypeptide consisting of a sequence of amino acid residues that is 80% identical in amino acid sequence to amino acid residue 114 to amino acid residue 246 of SEQ ID NO:2 comprising the sequence of SEQ ID NO:5; and c) a polypeptide consisting of a sequence of amino acid residues that is 80% identical in amino acid sequence to amino acid residue 51 to 246 of SEQ ID NO:2, the polypeptide consisting of Gly-Xaa-Xaa and Gly-Xaa-Pro repeats forming a collagen domain and comprising the sequence of SEQ ID NO:5.

Within another aspect is provided a fusion protein comprising a first portion and a second portion joined by a peptide bond, the first portion consisting of a polypeptide selected from the group consisting of: a) a polypeptide comprising a sequence of amino acid residues that is at least 75% identical in amino acid sequence to amino acid residue 51 to amino acid residue 246 of SEQ ID

NO:2; b) a polypeptide comprising a sequence of amino acid residues as shown in SEQ ID NO:2 from amino acid residue 1 to amino acid residue 246; c) a portion of the zacrp3 polypeptide of SEQ ID NO:2, comprising the collagen-like domain or a portion of the collagen-like domain capable of dimerization or oligomerization; d) a portion of the zacrp3 polypeptide of SEQ ID NO:2, comprising the Clq domain or an active portion of the Clq domain; or e) a portion of the zacrp3 polypeptide of SEQ ID NO:2 comprising of the collagen-like domain and the Clq domain; and the second portion comprising another polypeptide. Within one embodiment the first portion is selected from the group consisting of: a) a polypeptide consisting of the sequence of amino acid residue 51 to amino acid residue 113 of SEQ ID NO:2; b) a polypeptide consisting of the sequence of amino acid residue 114 to amino acid residue 246 of SEQ ID NO:2; c) a polypeptide consisting of the sequence of amino acid residue 51 to 246 of SEQ ID NO:2.

20 The invention also provides a polypeptide as described above; in combination with a pharmaceutically acceptable vehicle.

Within another aspect, the invention provides an antibody or antibody fragment that specifically binds to a polypeptide as described above. Within one embodiment the antibody is selected from the group consisting of: a) polyclonal antibody; b) murine monoclonal antibody; c) humanized antibody derived from b); and d) human monoclonal antibody. Within another embodiment the antibody fragment is selected from the group consisting of F(ab'), F(ab), Fab', Fab, Fv, scFv, and minimal recognition unit. Within another embodiment is provided an anti-idiotypic antibody that specifically binds to the antibody described above.

35 Within another aspect, the invention provides an isolated polynucleotide encoding a polypeptide comprising a sequence of amino acid residues that is at least 75%

identical in amino acid sequence to residues 51-246 of SEQ ID NO:2, wherein the sequence comprises: Gly-Xaa-Xaa or Gly-Xaa-Pro repeats forming a collagen domain, wherein Xaa is any amino acid; and a carboxyl-terminal C1q domain consisting of 10 beta strands. Within one embodiment the polypeptide is at least 90% identical in amino acid sequence to residues 23-246 of SEQ ID NO:2. Within another embodiment the collagen domain consists of 15 Gly-Xaa-Xaa repeats and 6 Gly-Xaa-Pro repeats. Within another embodiment the carboxyl-terminal C1q domain comprises the sequence of SEQ ID NO:5. Within another embodiment the carboxy-terminal C1q domain consists of amino acid residues 119-123, 140-142, 148-151, 155-158, 161-173, 175-182, 190-197, 200-212, 217-222 and 236-241 of SEQ ID NO:2. Within another embodiment any differences between the polypeptide and SEQ ID NO:2 are due to conservative amino acid substitutions. Within yet another embodiment the polypeptide specifically binds with an antibody that specifically binds with a polypeptide consisting of the amino acid sequence of SEQ ID NO:2. Within another embodiment the polypeptide comprises residues 23-246 of SEQ ID NO:2. Within another embodiment the collagen domain consists of amino acid residues 51-113 of SEQ ID NO:2. Within yet another embodiment the C1q domain consists of amino acid residues 114-246 of SEQ ID NO:2.

Also provided is an isolated polynucleotide selected from the group consisting of: a) a sequence of nucleotides from nucleotide 1 to nucleotide 1696 of SEQ ID NO:1; b) a sequence of nucleotides from nucleotide 69 to nucleotide 806 of SEQ ID NO:1; c) a sequence of nucleotides from nucleotide 135 to nucleotide 806 of SEQ ID NO:1; d) a sequence of nucleotides from nucleotide 219 to nucleotide 806 of SEQ ID NO:1; e) a sequence of nucleotides from nucleotide 408 to nucleotide 806 of SEQ ID NO:1; f) a sequence of nucleotides from nucleotide 69 to nucleotide 407 of SEQ ID NO:1; g) a sequence of nucleotides from nucleotide 135 to nucleotide 407 of SEQ

ID NO:1; h) a sequence of nucleotides from nucleotide 219 to nucleotide 407 of SEQ ID NO:1; i) a polynucleotide encoding a polypeptide, the polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 51 to 113 of SEQ ID NO:2; j) a polynucleotide encoding a polypeptide, the polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 114 to 246 of SEQ ID NO:2; k) a polynucleotide encoding a polypeptide, the polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 51 to 246 of SEQ ID NO:2; l) a polynucleotide encoding a polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 23 to 113 of SEQ ID NO:2; m) a polynucleotide that remains hybridized following stringent wash conditions to a polynucleotide consisting of the nucleotide sequence of SEQ ID NO:1, or the complement of SEQ ID NO:1; n) nucleotide sequences complementary to a), b), c), d), e), f), g), h), i), j), k), l) or m) and o) degenerate nucleotide sequences of i), j), k) or l).

Also provided is an isolated polynucleotide encoding a fusion protein comprises a first portion and a second portion joined by a peptide bond, the first portion is selected from the group consisting of: a) a polypeptide comprising a sequence of amino acid residues that is at least 75% identical in amino acid sequence to amino acid residues 51 to 246 of SEQ ID NO:2; b) a polypeptide comprising the sequence of amino acid residues 1 to 246 of SEQ ID NO:2; c) a polypeptide comprising the sequence of amino acid residues 23 to 246 of SEQ ID NO:2; d) a polypeptide comprising the sequence of amino acid residues 23 to 113 of SEQ ID NO:2; e) a polypeptide comprising the

sequence of amino acid residues 1 to 113 of SEQ ID NO:2;  
f) a portion of a polypeptide of SEQ ID NO:2 comprising  
the collagen-like domain or a portion of the collagen-like  
domain capable of dimerization or oligomerization; g) a  
5 portion of the polypeptide of SEQ ID NO:2 containing the  
Clq domain; or h) a portion of the polypeptide of SEQ ID  
NO:2 including the collagen-like domain and the Clq  
domain; and the second portion comprising another  
polypeptide.

10 Also provided is an isolated polynucleotide  
consisting of the sequence of nucleotide 1 to nucleotide  
738 of SEQ ID NO:10.

Within another aspect, the invention provides an  
expression vector comprising the following operably linked  
15 elements: a transcription promoter; a DNA segment encoding  
a polypeptide as described above; and a transcription  
terminator. Within one embodiment the DNA segment encodes  
a polypeptide that is at least 90% identical in amino acid  
sequence to residues 23-246 of SEQ ID NO:2. Within  
20 another embodiment the collagen domain consists of 15 Gly-  
Xaa-Xaa repeats and 6 Gly-Xaa-Pro repeats. Within another  
embodiment the carboxyl-terminal Clq domain comprises the  
sequence of SEQ ID NO:10. Within another embodiment the  
carboxy-terminal Clq domain consists of amino acid  
25 residues 119-123, 140-142, 148-151, 155-158, 161-173, 175-  
182, 190-197, 200-212, 217-222 and 236-241 of SEQ ID NO:2.  
Within another embodiment differences between the  
polypeptide and SEQ ID NO:2 are due to conservative amino  
acid substitutions. Within yet another embodiment the  
30 polypeptide specifically binds with an antibody that  
specifically binds with a polypeptide consisting of the  
amino acid sequence of SEQ ID NO:2. Within a further  
embodiment the DNA segment encodes a polypeptide  
comprising residues 23-246 of SEQ ID NO:2. Within another  
35 embodiment the collagen domain consists of amino acid  
residues 51-113 of SEQ ID NO:2. Within yet another  
embodiment the Clq domain consists of amino acid residues

114-246 of SEQ ID NO:2. Within yet another embodiment the DNA segment encodes a polypeptide covalently linked at the amino or carboxyl terminus to an affinity tag. Within another embodiment the DNA segment further encodes a secretory signal sequence operably linked to the polypeptide. Within a related embodiment the secretory signal sequence comprises residues 1-22 of SEQ ID NO:2.

Within another aspect, the invention provides a cultured cell into which has been introduced an expression vector as described above, wherein the cell expresses the polypeptide encoded by the DNA segment.

Within another aspect, the invention provides a method of producing a polypeptide comprising: culturing a cell into which has been introduced an expression vector as describe above; whereby the cell expresses the polypeptide encoded by the DNA segment; and recovering the expressed polypeptide.

#### BRIEF DESCRIPTION OF THE DRAWING

The Figure illustrates a multiple alignment of and zacrp3 polypeptide of the present invention and human ACRP30 (ACR3) (SEQ ID NO:3, Maeda et al., Biochem. Biophys. Res. Commun. 221:286-9, 1996) and human Clq C (SEQ ID NO:4, Sellar et al., Biochem J. 274:481-90, 1991 and Reid, Biochem J. 179:361-71, 1979). The multiple alignment performed using a Clustalx multiple alignment tool with the default settings: Blosum Series Weight Matricies, Gap Opening penalty:10.0, Gap Extension penalty:0.05. Multiple alignments were further hand tuned before computing percent identity.

#### DETAILED DESCRIPTION OF THE INVENTION

Prior to setting forth the invention in detail, it may be helpful to the understanding thereof to define the following terms.

The term "affinity tag" is used herein to denote a peptide segment that can be attached to a polypeptide to provide for purification or detection of the polypeptide or provide sites for attachment of the polypeptide to a substrate. In principal, any peptide or protein for which an antibody or other specific binding agent is available can be used as an affinity tag. Affinity tags include a poly-histidine tract, protein A (Nilsson et al., EMBO J. 4:1075, 1985; Nilsson et al., Methods Enzymol. 198:3, 1991), glutathione S transferase (Smith and Johnson, Gene 67:31, 1988), substance P, Flag™ peptide (Hopp et al., Biotechnology 6:1204-10, 1988; available from Eastman Kodak Co., New Haven, CT), streptavidin binding peptide, or other antigenic epitope or binding domain. See, in general Ford et al., Protein Expression and Purification 2: 95-107, 1991. DNAs encoding affinity tags are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ).

The term "allelic variant" denotes any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in phenotypic polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequence. The term allelic variant is also used herein to denote a protein encoded by an allelic variant of a gene.

The terms "amino-terminal" and "carboxyl-terminal" are used herein to denote positions within polypeptides and proteins. Where the context allows, these terms are used with reference to a particular sequence or portion of a polypeptide or protein to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a protein is located proximal to the carboxyl terminus of the reference sequence, but is not

necessarily at the carboxyl terminus of the complete protein.

The term "complement/anti-complement pair" denotes non-identical moieties that form a non-covalently associated, stable pair under appropriate conditions. For instance, biotin and avidin (or streptavidin) are prototypical members of a complement/anti-complement pair. Other exemplary complement/anti-complement pairs include receptor/ligand pairs, antibody/antigen (or hapten or epitope) pairs, sense/antisense polynucleotide pairs, and the like. Where subsequent dissociation of the complement/anti-complement pair is desirable, the complement/anti-complement pair preferably has a binding affinity of  $<10^9 \text{ M}^{-1}$ .

The term "complements of a polynucleotide molecule" is a polynucleotide molecule having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

The term "contig" denotes a polynucleotide that has a contiguous stretch of identical or complementary sequence to another polynucleotide. Contiguous sequences are said to "overlap" a given stretch of polynucleotide sequence either in their entirety or along a partial stretch of the polynucleotide. For example, representative contigs to the polynucleotide sequence 5'-ATGGCTTAGCTT-3' are 5'-TAGCTTgagtct-3' and 3'-gtcgacTACCGA-5'.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "expression vector" denotes a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments may include promoter and terminator sequences, and may optionally include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, and the like. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both.

The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, Nature 316:774-78, 1985).

An "isolated" polypeptide or protein is a polypeptide or protein that is found in a condition other than its native environment, such as apart from blood and animal tissue. In a preferred form, the isolated polypeptide is substantially free of other polypeptides, particularly other polypeptides of animal origin. It is preferred to provide the polypeptides in a highly purified form, i.e. greater than 95% pure, more preferably greater than 99% pure. When used in this context, the term "isolated" does not exclude the presence of the same polypeptide in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

The term "operably linked", when referring to DNA segments, denotes that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in the promoter and proceeds through the coding segment to the terminator.

The term "ortholog" denotes a polypeptide or protein obtained from one species that is the functional counterpart of a polypeptide or protein from a different species. Sequence differences among orthologs are the result of speciation.

"Paralogs" are distinct but structurally related proteins made by an organism. Paralogs are believed to arise through gene duplication. For example,  $\alpha$ -globin,  $\beta$ -globin, and myoglobin are paralogs of each other.

The term "polynucleotide" denotes a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules. Sizes of polynucleotides are expressed as base pairs (abbreviated "bp"), nucleotides ("nt"), or kilobases ("kb"). Where the context allows, the latter two terms may describe polynucleotides that are single-stranded or double-stranded. When the term is applied to double-stranded molecules it is used to denote overall length and will be understood to be equivalent to the term "base pairs". It will be recognized by those skilled in the art that the two strands of a double-stranded polynucleotide may differ slightly in length and that the ends thereof may be staggered as a result of enzymatic cleavage; thus all nucleotides within a double-stranded polynucleotide molecule may not be paired. Such unpaired ends will in general not exceed 20 nt in length.

A "polypeptide" is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than

about 10 amino acid residues are commonly referred to as "peptides".

"Probes and/or primers" as used herein can be RNA or DNA. DNA can be either cDNA or genomic DNA. 5 Polynucleotide probes and primers are single or double-stranded DNA or RNA, generally synthetic oligonucleotides, but may be generated from cloned cDNA or genomic sequences or its complements. Analytical probes will generally be at least 20 nucleotides in length, although somewhat 10 shorter probes (14-17 nucleotides) can be used. PCR primers are at least 5 nucleotides in length, preferably 15 or more nt, more preferably 20-30 nt. Short polynucleotides can be used when a small region of the gene is targeted for analysis. For gross analysis of 15 genes, a polynucleotide probe may comprise an entire exon or more. Probes can be labeled to provide a detectable signal, such as with an enzyme, biotin, a radionuclide, fluorophore, chemiluminescer, paramagnetic particle and the like, which are commercially available from many 20 sources, such as Molecular Probes, Inc., Eugene, OR, and Amersham Corp., Arlington Heights, IL, using techniques that are well known in the art.

The term "promoter" denotes a portion of a gene containing DNA sequences that provide for the binding of 25 RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

The term "receptor" denotes a cell-associated protein that binds to a bioactive molecule (i.e., a 30 ligand) and mediates the effect of the ligand on the cell. Membrane-bound receptors are characterized by a multi-domain structure comprising an extracellular ligand-binding domain and an intracellular effector domain that is typically involved in signal transduction. Binding of 35 ligand to receptor results in a conformational change in the receptor that causes an interaction between the effector domain and other molecule(s) in the cell. This

interaction in turn leads to an alteration in the metabolism of the cell. Metabolic events that are linked to receptor-ligand interactions include gene transcription, phosphorylation, dephosphorylation, increases in cyclic AMP production, mobilization of cellular calcium, mobilization of membrane lipids, cell adhesion, hydrolysis of inositol lipids and hydrolysis of phospholipids. Most nuclear receptors also exhibit a multi-domain structure, including an amino-terminal, transactivating domain, a DNA binding domain and a ligand binding domain. In general, receptors can be membrane bound, cytosolic or nuclear; monomeric (e.g., thyroid stimulating hormone receptor, beta-adrenergic receptor) or multimeric (e.g., PDGF receptor, growth hormone receptor, IL-3 receptor, GM-CSF receptor, G-CSF receptor, erythropoietin receptor and IL-6 receptor).

The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger peptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

A "soluble receptor" is a receptor polypeptide that is not bound to a cell membrane. Soluble receptors are most commonly ligand-binding receptor polypeptides that lack transmembrane and cytoplasmic domains. Soluble receptors can comprise additional amino acid residues, such as affinity tags that provide for purification of the polypeptide or provide sites for attachment of the polypeptide to a substrate, or immunoglobulin constant region sequences. Many cell-surface receptors have naturally occurring, soluble counterparts that are produced by proteolysis or translated from alternatively spliced mRNAs. Receptor polypeptides are said to be substantially free of transmembrane and intracellular polypeptide segments when they lack sufficient portions of

these segments to provide membrane anchoring or signal transduction, respectively.

The term "splice variant" is used herein to denote alternative forms of RNA transcribed from a gene. Splice variation arises naturally through use of alternative splicing sites within a transcribed RNA molecule, or less commonly between separately transcribed RNA molecules, and may result in several mRNAs transcribed from the same gene. Splice variants may encode polypeptides having altered amino acid sequence. The term splice variant is also used herein to denote a protein encoded by a splice variant of an mRNA transcribed from a gene.

Molecular weights and lengths of polymers determined by imprecise analytical methods (e.g., gel electrophoresis) will be understood to be approximate values. When such a value is expressed as "about" X or "approximately" X, the stated value of X will be understood to be accurate to  $\pm 10\%$ .

The present invention is based in part upon the discovery of a novel DNA sequence that encodes a polypeptide having homology to an adipocyte complement related protein (Acrp30). The novel DNA sequence encodes a polypeptide having an amino-terminal signal sequence, an adjacent N-terminal region of non-homology, a collagen domain composed of 21 Gly-Xaa-Xaa or Gly-Xaa-Pro repeats and a carboxy-terminal globular-like Clq domain followed by a long 3' untranslated region. The general polypeptide structure set forth above is shared by Acrp30 and Clq C. Other regions of homology, found in the carboxy-terminal globular Clq domain in the aligned proteins, are identified herein as useful primers for searching for other family members. Acrp30 and Clq C, for example, would be identified in a search using the primers. Intra-chain disulfide bonding may involve the cysteines at residues 39, 42 and 43 of SEQ ID NO:2.

The novel zacrp3 polypeptides of the present invention were initially identified by querying an EST database for homologs of ACRP30, characterized by a signal sequence, a collagen-like domain and a Clq domain.

5 Polypeptides corresponding to ESTs meeting those search criteria were compared to known sequences to identify proteins having homology to ACRP30. An assembled EST cluster was discovered and predicted to be a secreted protein. To identify the corresponding cDNA, a clone

10 considered likely to contain the entire coding sequence was used for sequencing. The resulting 1696 bp sequence is disclosed in SEQ ID NO:1. Comparison of the originally derived EST sequence with the sequence represented in SEQ ID NO:1 showed that there were two frame shifts and an

15 unspliced intron. The novel polypeptide encoded by the full length cDNA enabled the identification of a homolog relationship with adipocyte complement related protein Acrp30 (SEQ ID NO:3) and complement component Clq C (SEQ ID NO:4) as is shown in the Figure. Zacrp3 shares 27.5 and

20 25.7% identity at the amino acid level with human ACRP30 and Clq C respectfully. Clq C and ACRP30 share 32.4% identity. Within the Clq domain, zacrp3 shares 26.3 and 26% identity at the amino acid level when compared to human ACRP30 and Clq C respectfully. Clq C and ACRP30

25 share 38.2% identity over this region.

The full sequence of the zacrp3 polypeptide was obtained from a single clone believed to contain it, wherein the clone was obtained from a chest wall soft tissue library. This message was also found using

30 electronic searches, in libraries of connective tissues, digestive, skeletal, respiratory and nervous system tissues as well as urinary tract tissues.

The nucleotide sequence of zacrp3 is described in SEQ ID NO:1, and its deduced amino acid sequence is

35 described in SEQ ID NO:2. As described generally above, the zacrp3 polypeptide includes a signal sequence, ranging from amino acid 1 (Met) to amino acid residue 22 (Cys).

The mature polypeptide therefore ranges from amino acid 23 (Gln) to amino acid 246 (Lys). Within the mature polypeptide, an N-terminal region of no known homology is found, ranging between amino acid residue 23 (Gln) and 50 (Arg) of SEQ ID NO:2. In addition, a collagen-like domain is found between amino acid 51 (Gly) and 113 (Pro). In the collagen-like domain, 6 perfect Gly-Xaa-Pro and 15 imperfect Gly-Xaa-Xaa repeats are observed. Acrp30 contains 22 perfect or imperfect repeats. Proline residues found in this domain at amino acid residue 56, 59, 62 and 113 of SEQ ID NO:2 may be hydroxylated. The zacrp3 polypeptide also includes a carboxy-terminal Clq domain, ranging from about amino acid 114 (Pro) to 246 (Lys). There is a fair amount of conserved structure within the Clq domain to enable proper folding. An aromatic motif seen in all Clq domain containing proteins (F-X(5)-[ND]-X(4)-[FYWL]-X(6)-F-X(5)-G-X-Y-X-F-X-[FY] (SEQ ID NO:5) is found between residues 169 and 199 of SEQ ID NO:2. X represents any amino acid residue and the number in parentheses () indicates the amino acid number of residues. The amino acid residues contained within the square parentheses [] restrict the choice of amino acid residues at that particular position. Zacrp3 polypeptide, human Clq C and Acrp30 appear to be homologous within the collagen domain and in the Clq domain, but not in the N-terminal portion of the mature polypeptide.

Another aspect of the present invention includes zacrp3 polypeptide fragments. Preferred fragments include those containing the collagen-like domain of zacrp3 polypeptides, ranging from amino acid 1 (Met), 23 (Gln) or 51 (Gly) to amino acid 1113 (Pro) of SEQ ID NO:2, a portion of the zacrp3 polypeptide containing the collagen-like domain or a portion of the collagen-like domain capable of dimerization or oligomerization. As used herein the term "collagen" or "collagen-like domain" refers to a series of repeating triplet amino acid sequences, "repeats" or "collagen repeats" represented by

the motifs Gly-Xaa-Pro or Gly-Xaa-Xaa, where Xaa is any amino acid residue. Such domains may contain as many as 21 collagen repeats or more. Fragments or proteins containing such collagen-like domains may form homomeric constructs (dimers or oligomers of the same fragment or protein). Moreover, such fragments or proteins containing such collagen-like domains may form heteromeric constructs, usually trimers.

These fragments are particularly useful in the study of collagen dimerization or oligomerization or in formation of fusion proteins as described more fully below. Polynucleotides encoding such fragments are also encompassed by the present invention, including the group consisting of (a) polynucleotide molecule comprising a sequence of nucleotides as shown in SEQ ID NO:1 from nucleotide 1, 69, 135 or 219 to nucleotide 407; (b) polynucleotide molecules that encode a zacrp3 polypeptide fragment that is at least 80% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 51 (Gly) to amino acid residue 113 (Pro); (c) molecules complementary to (a) or (b); and (d) degenerate nucleotide sequences encoding a zacrp3 polypeptide collagen-like domain fragment.

Other preferred fragments include the globular C1q domain of zacrp3 polypeptides, ranging from amino acid 114 (Pro) to 246 (Lys) of SEQ ID NO:2, a portion of the zacrp3 polypeptide containing the C1q domain or an active portion of the C1q domain. Other C1q domain containing proteins include C1q A, B and C (Sellar et al., *ibid.*, Reid, *ibid.*, and Reid et al., *Biochem. J.* 203: 559-69, 1982), chipmunk hibernation-associated plasma proteins HP-20, HP-25 and HP-27 (Takamatsu et al., *Mol. Cell. Biol.* 13: 1516-21, 1993 and Kondo & Kondo, *J. Biol. Chem.* 267: 473-8, 1992), human precerebellin (Urade et al., *Proc. Natl. Acad. Sci. USA* 88:1069-73, 1991), human endothelial cell multimerin (Hayward et al., *J. Biol. Chem.* 270:18246-51, 1995) and vertebrate collagens type VIII and X

(Muragaki et al., Eur. J. Biochem. 197:615-22, 1991). The globular Clq domain of ACRP30 has been determined to have a 10 beta strand "jelly roll" topology (Shapiro and Scherer, Curr. Biol. 8:335-8, 1998) that shows significant  
5 homology to the TNF family and the zacrp3 sequence as represented by SEQ ID NO:2 contains all 10 beta-strands of this structure (amino acid residues 119-123, 140-142, 148-151, 155-158, 161-173, 175-182, 190-197, 200-212, 217-222 and 236-241 of SEQ ID NO:2). These strands have been  
10 designated "A", "A'", "B", "B'", "C", "D", "E", "F", "G" and "H" respectively.

Zacrp3 has two receptor binding loops, at amino acid residues 111-139 and 170-182, does zacrp3 have anything similar. The core receptor binding region is  
15 predicted to include amino acid residues 124-150 and 181-197 of SEQ ID NO:2. Amino acid residues 161 (Gly), 163 (Tyr), 212 (Leu) and 237 (Phe) appear to be conserved across the superfamily including CD40, TNF $\alpha$ , ACRP30 and zacrp3.

20 These fragments are particularly useful in the study or modulation of energy balance or neurotransmission, particularly diet- or stress-related neurotransmission. Anti-microbial activity may also be present in such fragments. The homology to TNF proteins  
25 suggests such fragments would be useful in obesity-related insulin resistance, immune regulation, inflammatory response, apoptosis and osteoclast maturation. Polynucleotides encoding such fragments are also encompassed by the present invention, including the group  
30 consisting of (a) polynucleotide molecules comprising a sequence of nucleotides as shown in SEQ ID NO:1 from nucleotide 408 to nucleotide 806; (b) polynucleotide molecules that encode a zacrp3 polypeptide fragment that is at least 80% identical to the amino acid sequence of  
35 SEQ ID NO:2 from amino acid residue 114 (Pro) to amino acid residue 246 (Lys); (c) molecules complementary to (a)

or (b); and (d) degenerate nucleotide sequences encoding a zacrp3 polypeptide Clq domain fragment.

Other zacrp3 polypeptide fragments of the present invention include both the collagen-like domain and the Clq domain ranging from amino acid residue 51 (Gly) to 806 (Lys) of SEQ ID NO:2. Polynucleotides encoding such fragments are also encompassed by the present invention, including the group consisting of (a) polynucleotide molecules comprising a sequence of nucleotides as shown in SEQ ID NO:1 from nucleotide 291 to nucleotide 806; (b) polynucleotide molecules that encode a zacrp3 polypeptide fragment that is at least 80% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 41 (Gly) to amino acid residue 246 (Lys); (c) molecules complementary to (a) or (b); and (d) degenerate nucleotide sequences encoding a zacrp3 polypeptide collagen-like domain-Clq domain fragment.

The highly conserved amino acids, particularly those in the carboxy-terminal Clq domain of the zacrp3 polypeptide, can be used as a tool to identify new family members. For instance, reverse transcription-polymerase chain reaction (RT-PCR) can be used to amplify sequences encoding the conserved motifs from RNA obtained from a variety of tissue sources. In particular, highly degenerate primers and their complements designed from conserved sequences are useful for this purpose. In particular, the following primers are useful for this purpose:

Amino acid residues 215-221 of SEQ ID NO:2  
GGN GAN SAR GTN TGG YT (SEQ ID NO:6)

Amino acid residues 160-165 of SEQ ID NO:2  
SN GNN NTN TAY TWY TTY R (SEQ ID NO:7)

Amino acid residues 237-242 of SEQ ID NO:2  
TTY DSN GGN TTY YTN HT (SEQ ID NO:8)

Amino acid residues 147-153 of SEQ ID NO:2

Y TWY RAY RBN WBN WSN GG (SEQ ID NO:9)

- 5 Probes corresponding to complements of the polynucleotides set forth above are also encompassed.

The present invention also provides polynucleotide molecules, including DNA and RNA molecules, that encode the zacrp3 polypeptides disclosed herein.

10 Those skilled in the art will readily recognize that, in view of the degeneracy of the genetic code, considerable sequence variation is possible among these polynucleotide molecules. SEQ ID NO:10 is a degenerate DNA sequence that encompasses all DNAs that encode the zacrp3 polypeptide of

15 SEQ ID NO:2. Those skilled in the art will recognize that the degenerate sequence of SEQ ID NO:10 also provides all RNA sequences encoding SEQ ID NO:2 by substituting U for T. Thus, zacrp3 polypeptide-encoding polynucleotides comprising nucleotide 1 to nucleotide 738 of SEQ ID NO:10

20 and their RNA equivalents are contemplated by the present invention. Table 1 sets forth the one-letter codes used within SEQ ID NO:10 to denote degenerate nucleotide positions. "Resolutions" are the nucleotides denoted by a code letter. "Complement" indicates the code for the

25 complementary nucleotide(s). For example, the code Y denotes either C or T, and its complement R denotes A or G, A being complementary to T, and G being complementary to C.

TABLE 1

Nucleotide	Resolution	Complement	Resolution
A	A	T	T
C	C	G	G
G	G	C	C
T	T	A	A
R	A G	Y	C T
Y	C T	R	A G
M	A C	K	G T
K	G T	M	A C
S	C G	S	C G
W	A T	W	A T
H	A C T	D	A G T
B	C G T	V	A C G
V	A C G	B	C G T
D	A G T	H	A C T
N	A C G T	N	A C G T

The degenerate codons used in SEQ ID NO:10,  
 5 encompassing all possible codons for a given amino acid,  
 are set forth in Table 2.

TABLE 2

Amino Acid	One Letter Code	Codons	Degenerate Codon
Cys	C	TGC TGT	TGY
Ser	S	AGC AGT TCA TCC TCG TCT	WSN
Thr	T	ACA ACC ACG ACT	ACN
Pro	P	CCA CCC CCG CCT	CCN
Ala	A	GCA GCC GCG GCT	GCN
Gly	G	GGA GGC GGG GGT	GGN
Asn	N	AAC AAT	AAY
Asp	D	GAC GAT	GAY
Glu	E	GAA GAG	GAR
Gln	Q	CAA CAG	CAR
His	H	CAC CAT	CAY
Arg	R	AGA AGG CGA CGC CGG CGT	MGN
Lys	K	AAA AAG	AAR
Met	M	ATG	ATG
Ile	I	ATA ATC ATT	ATH
Leu	L	CTA CTC CTG CTT TTA TTG	YTN
Val	V	GTA GTC GTG GTT	GTN
Phe	F	TTC TTT	TTY
Tyr	Y	TAC TAT	TAY
Trp	W	TGG	TGG
Ter	.	TAA TAG TGA	TRR
Asn Asp	B		RAY
Glu Gln	Z		SAR
Any	X		NNN

One of ordinary skill in the art will appreciate that some ambiguity is introduced in determining a degenerate codon, representative of all possible codons encoding each amino acid. For example, the degenerate  
5 codon for serine (WSN) can, in some circumstances, encode arginine (AGR), and the degenerate codon for arginine (MGN) can, in some circumstances, encode serine (AGY). A similar relationship exists between codons encoding phenylalanine and leucine. Thus, some polynucleotides  
10 encompassed by the degenerate sequence may encode variant amino acid sequences, but one of ordinary skill in the art can easily identify such variant sequences by reference to the amino acid sequence of SEQ ID NO:2. Variant sequences can be readily tested for functionality as described  
15 herein.

One of ordinary skill in the art will also appreciate that different species can exhibit "preferential codon usage." In general, see, Grantham, et al., Nuc. Acids Res. 8:1893-912, 1980; Haas, et al. Curr.  
20 Biol. 6:315-24, 1996; Wain-Hobson, et al., Gene 13:355-64, 1981; Grosjean and Fiers, Gene 18:199-209, 1982; Holm, Nuc. Acids Res. 14:3075-87, 1986; Ikemura, J. Mol. Biol. 158:573-97, 1982. As used herein, the term "preferential codon usage" or "preferential codons" is a term of art  
25 referring to protein translation codons that are most frequently used in cells of a certain species, thus favoring one or a few representatives of the possible codons encoding each amino acid (See Table 2). For example, the amino acid threonine (Thr) may be encoded by  
30 ACA, ACC, ACG, or ACT, but in mammalian cells ACC is the most commonly used codon; in other species, for example, insect cells, yeast, viruses or bacteria, different Thr codons may be preferential. Preferential codons for a particular species can be introduced into the  
35 polynucleotides of the present invention by a variety of methods known in the art. Introduction of preferential codon sequences into recombinant DNA can, for example,

enhance production of the protein by making protein translation more efficient within a particular cell type or species. Therefore, the degenerate codon sequence disclosed in SEQ ID NO:10 serves as a template for  
5 optimizing expression of polynucleotides in various cell types and species commonly used in the art and disclosed herein. Sequences containing preferential codons can be tested and optimized for expression in various species, and tested for functionality as disclosed herein.

10 The present invention further provides variant polypeptides and nucleic acid molecules that represent counterparts from other species (orthologs). These species include, but are not limited to mammalian, avian, amphibian, reptile, fish, insect and other vertebrate and  
15 invertebrate species. Of particular interest are zacrp3 polypeptides from other mammalian species, including murine, porcine, ovine, bovine, canine, feline, equine, and other primate polypeptides. A murine zacrp2 homolog (SEQ ID NO:12) has been identified. The polynucleotide  
20 sequence encoding this murine zacrp2 polypeptide disclosed in SEQ ID NO:11. Orthologs of human zacrp3 can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a cDNA can be cloned using mRNA  
25 obtained from a tissue or cell type that expresses zacrp3 as disclosed herein. Suitable sources of mRNA can be identified by probing northern blots with probes designed from the sequences disclosed herein. A library is then prepared from mRNA of a positive tissue or cell line.

30 An zacrp3-encoding cDNA can then be isolated by a variety of methods, such as by probing with a complete or partial human cDNA or with one or more sets of degenerate probes based on the disclosed sequences. A cDNA can also be cloned using the polymerase chain  
35 reaction with primers designed from the representative human zacrp3 sequences disclosed herein. Within an additional method, the cDNA library can be used to

transform or transfect host cells, and expression of the cDNA of interest can be detected with an antibody to zacrp3 polypeptide. Similar techniques can also be applied to the isolation of genomic clones.

5 Those skilled in the art will recognize that the sequence disclosed in SEQ ID NO:1 represents a single allele of human zacrp3, and that allelic variation and alternative splicing are expected to occur. Allelic variants of this sequence can be cloned by probing cDNA or  
10 genomic libraries from different individuals according to standard procedures. Allelic variants of the nucleotide sequence shown in SEQ ID NO:1, including those containing silent mutations and those in which mutations result in amino acid sequence changes, are within the scope of the  
15 present invention, as are proteins which are allelic variants of SEQ ID NO:2. cDNA molecules generated from alternatively spliced mRNAs, which retain the properties of the zacrp3 polypeptide are included within the scope of the present invention, as are polypeptides encoded by such  
20 cDNAs and mRNAs. Allelic variants and splice variants of these sequences can be cloned by probing cDNA or genomic libraries from different individuals or tissues according to standard procedures known in the art.

Within preferred embodiments of the invention,  
25 the isolated nucleic acid molecules can hybridize under stringent conditions to nucleic acid molecules having the nucleotide sequence of SEQ ID NO:1 or to nucleic acid molecules having a nucleotide sequence complementary to SEQ ID NO:1. In general, stringent conditions are  
30 selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe.

35 A pair of nucleic acid molecules, such as DNA-DNA, RNA-RNA and DNA-RNA, can hybridize if the nucleotide sequences have some degree of complementarity. Hybrids can

tolerate mismatched base pairs in the double helix, but the stability of the hybrid is influenced by the degree of mismatch. The  $T_m$  of the mismatched hybrid decreases by 1°C for every 1-1.5% base pair mismatch. Varying the stringency of the hybridization conditions allows control over the degree of mismatch that will be present in the hybrid. The degree of stringency increases as the hybridization temperature increases and the ionic strength of the hybridization buffer decreases. Stringent hybridization conditions encompass temperatures of about 5-25°C below the  $T_m$  of the hybrid and a hybridization buffer having up to 1 M Na<sup>+</sup>. Higher degrees of stringency at lower temperatures can be achieved with the addition of formamide which reduces the  $T_m$  of the hybrid about 1°C for each 1% formamide in the buffer solution. Generally, such stringent conditions include temperatures of 20-70°C and a hybridization buffer containing up to 6x SSC and 0-50% formamide. A higher degree of stringency can be achieved at temperatures of from 40-70°C with a hybridization buffer having up to 4x SSC and from 0-50% formamide. Highly stringent conditions typically encompass temperatures of 42-70°C with a hybridization buffer having up to 1x SSC and 0-50% formamide. Different degrees of stringency can be used during hybridization and washing to achieve maximum specific binding to the target sequence. Typically, the washes following hybridization are performed at increasing degrees of stringency to remove non-hybridized polynucleotide probes from hybridized complexes.

The above conditions are meant to serve as a guide and it is well within the abilities of one skilled in the art to adapt these conditions for use with a particular polypeptide hybrid. The  $T_m$  for a specific target sequence is the temperature (under defined conditions) at which 50% of the target sequence will hybridize to a perfectly matched probe sequence. Those conditions which influence the  $T_m$  include, the size and

base pair content of the polynucleotide probe, the ionic strength of the hybridization solution, and the presence of destabilizing agents in the hybridization solution. Numerous equations for calculating  $T_m$  are known in the art, and are specific for DNA, RNA and DNA-RNA hybrids and polynucleotide probe sequences of varying length (see, for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition (Cold Spring Harbor Press 1989); Ausubel et al., (eds.), Current Protocols in Molecular Biology (John Wiley and Sons, Inc. 1987); Berger and Kimmel (eds.), Guide to Molecular Cloning Techniques, (Academic Press, Inc. 1987); and Wetmur, Crit. Rev. Biochem. Mol. Biol. 26:227 (1990)). Sequence analysis software, such as OLIGO 6.0 (LSR; Long Lake, MN) and Primer Premier 4.0 (Premier Biosoft International; Palo Alto, CA), as well as sites on the Internet, are available tools for analyzing a given sequence and calculating  $T_m$  based on user defined criteria. Such programs can also analyze a given sequence under defined conditions and identify suitable probe sequences. Typically, hybridization of longer polynucleotide sequences, >50 base pairs, is performed at temperatures of about 20-25°C below the calculated  $T_m$ . For smaller probes, <50 base pairs, hybridization is typically carried out at the  $T_m$  or 5-10°C below. This allows for the maximum rate of hybridization for DNA-DNA and DNA-RNA hybrids.

The length of the polynucleotide sequence influences the rate and stability of hybrid formation. Smaller probe sequences, <50 base pairs, reach equilibrium with complementary sequences rapidly, but may form less stable hybrids. Incubation times of anywhere from minutes to hours can be used to achieve hybrid formation. Longer probe sequences come to equilibrium more slowly, but form more stable complexes even at lower temperatures. Incubations are allowed to proceed overnight or longer. Generally, incubations are carried out for a period equal to three times the calculated Cot time. Cot time, the

time it takes for the polynucleotide sequences to reassociate, can be calculated for a particular sequence by methods known in the art.

The base pair composition of polynucleotide  
5 sequence will effect the thermal stability of the hybrid complex, thereby influencing the choice of hybridization temperature and the ionic strength of the hybridization buffer. A-T pairs are less stable than G-C pairs in aqueous solutions containing sodium chloride. Therefore,  
10 the higher the G-C content, the more stable the hybrid. Even distribution of G and C residues within the sequence also contribute positively to hybrid stability. In addition, the base pair composition can be manipulated to alter the  $T_m$  of a given sequence. For example, 5-  
15 methyldeoxycytidine can be substituted for deoxycytidine and 5-bromodeoxuridine can be substituted for thymidine to increase the  $T_m$ , whereas 7-deaz-2'-deoxyguanosine can be substituted for guanosine to reduce dependence on  $T_m$ .

The ionic concentration of the hybridization  
20 buffer also affects the stability of the hybrid. Hybridization buffers generally contain blocking agents such as Denhardt's solution (Sigma Chemical Co., St. Louis, Mo.), denatured salmon sperm DNA, tRNA, milk powders (BLOTTO), heparin or SDS, and a  $Na^+$  source, such as  
25 SSC (1x SSC: 0.15 M sodium chloride, 15 mM sodium citrate) or SSPE (1x SSPE: 1.8 M NaCl, 10 mM  $NaH_2PO_4$ , 1 mM EDTA, pH 7.7). By decreasing the ionic concentration of the buffer, the stability of the hybrid is increased. Typically, hybridization buffers contain from between 10  
30 mM - 1 M  $Na^+$ . The addition of destabilizing or denaturing agents such as formamide, tetralkylammonium salts, guanidinium cations or thiocyanate cations to the hybridization solution will alter the  $T_m$  of a hybrid. Typically, formamide is used at a concentration of up to  
35 50% to allow incubations to be carried out at more convenient and lower temperatures. Formamide also acts to reduce non-specific background when using RNA probes.

As an illustration, a nucleic acid molecule encoding a variant zacrp3 polypeptide can be hybridized with a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1 (or its complement) at 42°C overnight in a solution comprising 50% formamide, 5x SSC (1x SSC: 0.15 M sodium chloride and 15 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution (100x Denhardt's solution: 2% (w/v) Ficoll 400, 2% (w/v) polyvinylpyrrolidone, and 2% (w/v) bovine serum albumin), 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA. One of skill in the art can devise variations of these hybridization conditions. For example, the hybridization mixture can be incubated at a higher or lower temperature, such as about 65°C, in a solution that does not contain formamide. Moreover, premixed hybridization solutions are available (e.g., EXPRESSHYB Hybridization Solution from CLONTECH Laboratories, Inc.), and hybridization can be performed according to the manufacturer's instructions.

Following hybridization, the nucleic acid molecules can be washed to remove non-hybridized nucleic acid molecules under stringent conditions, or under highly stringent conditions. Typical stringent washing conditions include washing in a solution of 0.5x-2x SSC with 0.1% sodium dodecyl sulfate (SDS) at 55-65°C. That is, nucleic acid molecules encoding a variant zacrp3 polypeptide hybridize with a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1 (or its complement) under stringent washing conditions, in which the wash stringency is equivalent to 0.5x-2x SSC with 0.1% SDS at 50-65°C, including 0.5x SSC with 0.1% SDS at 55°C, or 2x SSC with 0.1% SDS at 65°C. One of skill in the art can readily devise equivalent conditions, for example, by substituting SSPE for SSC in the wash solution.

Typical highly stringent washing conditions include washing in a solution of 0.1x-0.2x SSC with 0.1% sodium dodecyl sulfate (SDS) at 50-65°C. In other words,

nucleic acid molecules encoding a variant zacrp3 polypeptide hybridize with a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1 (or its complement) under highly stringent washing conditions, in which the  
5 wash stringency is equivalent to 0.1x-0.2x SSC with 0.1% SDS at 50-65°C, including 0.1x SSC with 0.1% SDS at 50°C, or 0.2x SSC with 0.1% SDS at 65°C.

The present invention also provides isolated zacrp3 polypeptides that have a substantially similar  
10 sequence identity to the polypeptides of SEQ ID NO:2, or their orthologs. The term "substantially similar sequence identity" is used herein to denote polypeptides having at least 70%, at least 80%, at least 90%, at least 95% or greater than 95% sequence identity to the sequences shown  
15 in SEQ ID NO:2, or their orthologs. The present invention also includes polypeptides that comprise an amino acid sequence having at least 70%, at least 80%, at least 90%, at least 95% or greater than 95% sequence identity to the sequence of amino acid residues 51 to 246 of SEQ ID NO:2.  
20 The present invention further includes nucleic acid molecules that encode such polypeptides. Methods for determining percent identity are described below.

The present invention also contemplates zacrp3 variant nucleic acid molecules that can be identified  
25 using two criteria: a determination of the similarity between the encoded polypeptide with the amino acid sequence of SEQ ID NO:2, and a hybridization assay, as described above. Such zacrp3 variants include nucleic acid molecules (1) that hybridize with a nucleic acid  
30 molecule having the nucleotide sequence of SEQ ID NO:1 (or its complement) under stringent washing conditions, in which the wash stringency is equivalent to 0.5x-2x SSC with 0.1% SDS at 50-65°C, and (2) that encode a polypeptide having at least 70%, at least 80%, at least  
35 90%, at least 95% or greater than 95% sequence identity to the amino acid sequence of SEQ ID NO:2. Alternatively, zacrp3 variants can be characterized as nucleic acid

molecules (1) that hybridize with a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1 (or its complement) under highly stringent washing conditions, in which the wash stringency is equivalent to 0.1x-0.2x SSC with 0.1% SDS at 50-65°C, and (2) that encode a polypeptide having at least 70%, at least 80%, at least 90%, at least 95% or greater than 95% sequence identity to the amino acid sequence of SEQ ID NO:2.

Percent sequence identity is determined by conventional methods. See, for example, Altschul et al., Bull. Math. Bio. 48:603, 1986, and Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA 89:10915, 1992. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "BLOSUM62" scoring matrix of Henikoff and Henikoff (ibid.) as shown in Table 3 (amino acids are indicated by the standard one-letter codes). The percent identity is then calculated as: ([Total number of identical matches]/ [length of the longer sequence plus the number of gaps introduced into the longer sequence in order to align the two sequences])(100).

Table 3

	A	R	N	D	C	Q	E	G	H	I	L	K	M	F	P	S	T	W	Y	V	
5	A	4																			
	R	-1	5																		
	N	-2	0	6																	
	D	-2	-2	1	6																
	C	0	-3	-3	-3	9															
10	Q	-1	1	0	0	-3	5														
	E	-1	0	0	2	-4	2	5													
	G	0	-2	0	-1	-3	-2	-2	6												
	H	-2	0	1	-1	-3	0	0	-2	8											
	I	-1	-3	-3	-3	-1	-3	-3	-4	-3	4										
	L	-1	-2	-3	-4	-1	-2	-3	-4	-3	2	4									
15	K	-1	2	0	-1	-3	1	1	-2	-1	-3	-2	5								
	M	-1	-1	-2	-3	-1	0	-2	-3	-2	1	2	-1	5							
	F	-2	-3	-3	-3	-2	-3	-3	-3	-1	0	0	-3	0	6						
	P	-1	-2	-2	-1	-3	-1	-1	-2	-2	-3	-3	-1	-2	-4	7					
	S	1	-1	1	0	-1	0	0	0	-1	-2	-2	0	-1	-2	-1	4				
20	T	0	-1	0	-1	-1	-1	-1	-2	-2	-1	-1	-1	-2	-1	1	5				
	W	-3	-3	-4	-4	-2	-2	-3	-2	-2	-3	-2	-3	-1	1	-4	-3	-2	11		
	Y	-2	-2	-2	-3	-2	-1	-2	-3	2	-1	-1	-2	-1	3	-3	-2	-2	2	7	
	V	0	-3	-3	-3	-1	-2	-2	-3	-3	3	1	-2	1	-1	-2	-2	0	-3	-1	4

Those skilled in the art appreciate that there are many established algorithms available to align two amino acid sequences. The "FASTA" similarity search algorithm of Pearson and Lipman is a suitable protein alignment method for examining the level of identity shared by an amino acid sequence disclosed herein and the amino acid sequence of a putative variant zacrp3. The FASTA algorithm is described by Pearson and Lipman, Proc. Nat. Acad. Sci. USA 85:2444, 1988, and by Pearson, Meth. Enzymol. 183:63, 1990.

Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO:2) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then re-scored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence and the ktup value), then the trimmed initial regions are examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman and Wunsch, J. Mol. Biol. 48:444, 1970; Sellers, SIAM J. Appl. Math. 26:787, 1974), which allows for amino acid insertions and deletions. Illustrative parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as

explained in Appendix 2 of Pearson, Meth. Enzymol. 183:63, 1990.

FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from four to six.

The present invention includes nucleic acid molecules that encode a polypeptide having one or more "conservative amino acid substitutions," compared with the amino acid sequence of SEQ ID NO:2. Conservative amino acid substitutions can be based upon the chemical properties of the amino acids. That is, variants can be obtained that contain one or more amino acid substitutions of SEQ ID NO:2, in which an alkyl amino acid is substituted for an alkyl amino acid in a zacrp3 amino acid sequence, an aromatic amino acid is substituted for an aromatic amino acid in a zacrp3 amino acid sequence, a sulfur-containing amino acid is substituted for a sulfur-containing amino acid in a zacrp3 amino acid sequence, a hydroxy-containing amino acid is substituted for a hydroxy-containing amino acid in a zacrp3 amino acid sequence, an acidic amino acid is substituted for an acidic amino acid in a zacrp3 amino acid sequence, a basic amino acid is substituted for a basic amino acid in a zacrp3 amino acid sequence, or a dibasic monocarboxylic amino acid is substituted for a dibasic monocarboxylic amino acid in a zacrp3 amino acid sequence.

Among the common amino acids, for example, a "conservative amino acid substitution" is illustrated by a substitution among amino acids within each of the following groups: (1) glycine, alanine, valine, leucine, and isoleucine, (2) phenylalanine, tyrosine, and tryptophan, (3) serine and threonine, (4) aspartate and glutamate, (5) glutamine and asparagine, and (6) lysine, arginine and histidine.

The BLOSUM62 table is an amino acid substitution matrix derived from about 2,000 local multiple alignments of protein sequence segments, representing highly conserved regions of more than 500 groups of related proteins (Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA 89:10915, 1992). Accordingly, the BLOSUM62 substitution frequencies can be used to define conservative amino acid substitutions that may be introduced into the amino acid sequences of the present invention. Although it is possible to design amino acid substitutions based solely upon chemical properties (as discussed above), the language "conservative amino acid substitution" preferably refers to a substitution represented by a BLOSUM62 value of greater than -1. For example, an amino acid substitution is conservative if the substitution is characterized by a BLOSUM62 value of 0, 1, 2, or 3. According to this system, preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 1 (e.g., 1, 2 or 3), while more preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 2 (e.g., 2 or 3).

Conservative amino acid changes in a zacrp3 gene can be introduced by substituting nucleotides for the nucleotides recited in SEQ ID NO:1. Such "conservative amino acid" variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reaction, and the like (see Ausubel (1995) at pages 8-10 to 8-22; and McPherson (ed.), Directed Mutagenesis: A Practical Approach (IRL Press 1991)). The ability of such variants to promote the energy balance modulating or other properties of the wild-type protein can be determined using a standard methods, such as the assays described herein. Alternatively, a variant zacrp3 polypeptide can be identified by the ability to specifically bind anti-zacrp3 antibodies.

The proteins of the present invention can also comprise non-naturally occurring amino acid residues. Non-naturally occurring amino acids include, without limitation, *trans*-3-methylproline, 2,4-methanoproline, 5 *cis*-4-hydroxyproline, *trans*-4-hydroxyproline, *N*-methylglycine, *allo*-threonine, methylthreonine, hydroxyethylcysteine, hydroxyethylhomocysteine, nitroglutamine, homoglutamine, pipecolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, 3,3-dimethylproline, 10 *tert*-leucine, norvaline, 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, and 4-fluorophenylalanine. Several methods are known in the art for incorporating non-naturally occurring amino acid residues into proteins. For example, an *in vitro* system can be 15 employed wherein nonsense mutations are suppressed using chemically aminoacylated suppressor tRNAs. Methods for synthesizing amino acids and aminoacylating tRNA are known in the art. Transcription and translation of plasmids containing nonsense mutations is typically carried out in 20 a cell-free system comprising an *E. coli* S30 extract and commercially available enzymes and other reagents. Proteins are purified by chromatography. See, for example, Robertson et al., J. Am. Chem. Soc. 113:2722, 1991, Ellman et al., Methods Enzymol. 202:301, 1991, Chung et al., 25 Science 259:806, 1993, and Chung et al., Proc. Nat. Acad. Sci. USA 90:10145, 1993.

In a second method, translation is carried out in Xenopus oocytes by microinjection of mutated mRNA and chemically aminoacylated suppressor tRNAs (Turcatti et 30 al., J. Biol. Chem. 271:19991, 1996). Within a third method, E. coli cells are cultured in the absence of a natural amino acid that is to be replaced (e.g., phenylalanine) and in the presence of the desired non-naturally occurring amino acid(s) (e.g., 2- 35 azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, or 4-fluorophenylalanine). The non-naturally occurring amino acid is incorporated into the protein in place of

its natural counterpart. See, Koide et al., Biochem. 33:7470, 1994. Naturally occurring amino acid residues can be converted to non-naturally occurring species by in vitro chemical modification. Chemical modification can be  
5 combined with site-directed mutagenesis to further expand the range of substitutions (Wynn and Richards, Protein Sci. 2:395, 1993).

A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic  
10 code, non-naturally occurring amino acids, and unnatural amino acids may be substituted for zacrp3 amino acid residues.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis and  
15 screening, such as those disclosed by Reidhaar-Olson and Sauer (Science 241:53, 1988) or Bowie and Sauer (Proc. Nat. Acad. Sci. USA 86:2152, 1989). Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, selecting for  
20 functional polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., Biochem. 30:10832, 1991, Ladner et al., U.S. Patent  
25 No. 5,223,409, Huse, international publication No. WO 92/06204, and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986, and Ner et al., DNA 7:127, 1988).

Variants of the disclosed zacrp3 nucleotide and polypeptide sequences can also be generated through DNA  
30 shuffling as disclosed by Stemmer, Nature 370:389, 1994, Stemmer, Proc. Nat. Acad. Sci. USA 91:10747, 1994, and international publication No. WO 97/20078. Briefly, variant DNA molecules are generated by in vitro homologous recombination by random fragmentation of a parent DNA  
35 followed by reassembly using PCR, resulting in randomly introduced point mutations. This technique can be modified by using a family of parent DNA molecules, such

as allelic variants or DNA molecules from different species, to introduce additional variability into the process. Selection or screening for the desired activity, followed by additional iterations of mutagenesis and assay  
5 provides for rapid "evolution" of sequences by selecting for desirable mutations while simultaneously selecting against detrimental changes.

Mutagenesis methods as disclosed herein can be combined with high-throughput, automated screening methods  
10 to detect activity of cloned, mutagenized polypeptides in host cells. Mutagenized DNA molecules that encode biologically active polypeptides, or polypeptides that bind with anti-zacrp3 antibodies, can be recovered from the host cells and rapidly sequenced using modern  
15 equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

Essential amino acids in the polypeptides of the  
20 present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244:1081, 1989, Bass et al., Proc. Nat. Acad. Sci. USA 88:4498, 1991, Coombs and Corey, "Site-  
25 Directed Mutagenesis and Protein Engineering," in Proteins: Analysis and Design, Angeletti (ed.), pages 259-311 (Academic Press, Inc. 1998)). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant  
30 molecules are tested for biological activity as disclosed below to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., J. Biol. Chem. 271:4699, 1996. The identities of essential amino acids can also be inferred from analysis of  
35 homologies with zacrp3.

The location of zacrp3 receptor binding domains can be identified by physical analysis of structure, as

determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., Science 255:306, 1992, Smith et al., J. Mol. Biol. 224:899, 1992, and Wlodaver et al., FEBS Lett. 309:59, 1992. Moreover, zacrp3 labeled with biotin or FITC can be used for expression cloning of zacrp3 receptors.

10           The present invention also provides polypeptide fragments or peptides comprising an epitope-bearing portion of a zacrp3 polypeptide described herein. Such fragments or peptides may comprise an "immunogenic epitope," which is a part of a protein that elicits an antibody response when the entire protein is used as an immunogen. Immunogenic epitope-bearing peptides can be identified using standard methods (see, for example, Geysen et al., Proc. Nat. Acad. Sci. USA 81:3998, 1983).

20           In contrast, polypeptide fragments or peptides may comprise an "antigenic epitope," which is a region of a protein molecule to which an antibody can specifically bind. Certain epitopes consist of a linear or contiguous stretch of amino acids, and the antigenicity of such an epitope is not disrupted by denaturing agents. It is known in the art that relatively short synthetic peptides that can mimic epitopes of a protein can be used to stimulate the production of antibodies against the protein (see, for example, Sutcliffe et al., Science 219:660, 1983). Accordingly, antigenic epitope-bearing peptides and polypeptides of the present invention are useful to raise antibodies that bind with the polypeptides described herein.

35           Antigenic epitope-bearing peptides and polypeptides preferably contain at least four to ten amino acids, at least ten to fifteen amino acids, or about 15 to about 30 amino acids of SEQ ID NO:2. Such epitope-bearing peptides and polypeptides can be produced by fragmenting a

zacrp3 polypeptide, or by chemical peptide synthesis, as described herein. Moreover, epitopes can be selected by phage display of random peptide libraries (see, for example, Lane and Stephen, Curr. Opin. Immunol. 5:268, 1993, and Cortese et al., Curr. Opin. Biotechnol. 7:616, 1996). Standard methods for identifying epitopes and producing antibodies from small peptides that comprise an epitope are described, for example, by Mole, "Epitope Mapping," in Methods in Molecular Biology, Vol. 10, Manson (ed.), pages 105-16 (The Humana Press, Inc. 1992), Price, "Production and Characterization of Synthetic Peptide-Derived Antibodies," in Monoclonal Antibodies: Production, Engineering, and Clinical Application, Ritter and Ladyman (eds.), pages 60-84 (Cambridge University Press 1995), and Coligan et al. (eds.), Current Protocols in Immunology, pages 9.3.1 - 9.3.5 and pages 9.4.1 - 9.4.11 (John Wiley & Sons 1997).

Regardless of the particular nucleotide sequence of a variant zacrp3 gene, the gene encodes a polypeptide that is characterized by its energy balance modulating activity or other activities of the wild-type protein, or by the ability to bind specifically to an anti-zacrp3 antibody. More specifically, variant zacrp3 genes encode polypeptides which exhibit at least 50%, and preferably, greater than 70, 80, or 90%, of the activity of polypeptide encoded by the human zacrp3 gene described herein.

For any zacrp3 polypeptide, including variants and fusion proteins, one of ordinary skill in the art can readily generate a fully degenerate polynucleotide sequence encoding that variant using the information set forth in Tables 1 and 2 above. Moreover, those of skill in the art can use standard software to devise zacrp3 variants based upon the nucleotide and amino acid sequences described herein. Accordingly, the present invention includes a computer-readable medium encoded with a data structure that provides at least one of the

following sequences: SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:10. Suitable forms of computer-readable media include magnetic media and optically-readable media. Examples of magnetic media include a hard or fixed drive, a random access memory (RAM) chip, a floppy disk, digital linear tape (DLT), a disk cache, and a ZIP disk. Optically readable media are exemplified by compact discs (e.g., CD-read only memory (ROM), CD-rewritable (RW), and CD-recordable), and digital versatile/video discs (DVD) (e.g., DVD-ROM, DVD-RAM, and DVD+RW).

The present invention also provides zacrp3 fusion proteins. For example, fusion proteins of the present invention encompass (1) a polypeptide selected from the group consisting of: (a) polypeptide molecules comprising a sequence of amino acid residues as shown in SEQ ID NO:2 from amino acid residue 1 (Met), 23 (Gln) or 51 (Gly) to amino acid residue 246 (Lys); (b) polypeptide molecules ranging from amino acid 51 (Gly) to amino acid 113 (Pro) of SEQ ID NO:2, a portion of the zacrp3 polypeptide containing the collagen-like domain or a portion of the collagen-like domain capable of dimerization or oligomerization; (c) polypeptide molecules ranging from amino acid 114 (Pro) to 246 (Lys) of SEQ ID NO:2, a portion of the zacrp3 polypeptide containing the Clq domain or an active portion of the Clq domain; or (d) polypeptide molecules ranging from amino acid 51 (Gly) to 246 (Lys), a portion of the zacrp3 polypeptide including the collagen-like domain and the Clq domain; and (2) another polypeptide. The other polypeptide may be alternative or additional Clq domain, an alternative or additional collagen-like domain, a signal peptide to facilitate secretion of the fusion protein or the like. The globular domain of complement binds IgG, thus, the globular domain of zacrp3 polypeptide, fragment or fusion may have a similar role.

Zacrp3 polypeptides, ranging from amino acid 1 (Met) to amino acid 246 (Lys); the mature zacrp3

polypeptides, ranging from amino acid 23 (Gln) to amino acid 246 (Lys); or the secretion leader fragments thereof, which fragments range from amino acid 1 (Met) to amino acid 22 (Cys) may be used in the study of secretion of proteins from cells. In preferred embodiments of this aspect of the present invention, the mature polypeptides are formed as fusion proteins with putative secretory signal sequences; plasmids bearing regulatory regions capable of directing the expression of the fusion protein is introduced into test cells; and secretion of mature protein is monitored. The monitoring may be done by techniques known in the art, such as HPLC and the like.

The polypeptides of the present invention, including full-length proteins, fragments thereof and fusion proteins, can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells. Eukaryotic cells, particularly cultured cells of multicellular organisms, are preferred. Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., *ibid.*, and Ausubel et al. *ibid.*

In general, a DNA sequence encoding a zacrp3 polypeptide of the present invention is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator within an expression vector. The vector will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors and

other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers.

5           To direct a zacrp3 polypeptide into the secretory pathway of a host cell, a secretory signal sequence (also known as a leader sequence, signal sequence, prepro sequence or pre-sequence) is provided in the expression vector. The secretory signal sequence may  
10 be that of the zacrp3 polypeptide, or may be derived from another secreted protein (e.g., t-PA) or synthesized de novo. The secretory signal sequence is joined to the zacrp3 polypeptide DNA sequence in the correct reading frame. Secretory signal sequences are commonly positioned  
15 5' to the DNA sequence encoding the polypeptide of interest, although certain signal sequences may be positioned elsewhere in the DNA sequence of interest (see, e.g., Welch et al., U.S. Patent No. 5,037,743; Holland et al., U.S. Patent No. 5,143,830). Conversely, the signal  
20 sequence portion of the zacrp3 polypeptide (amino acid residues 1-22 of SEQ ID NO:2) may be employed to direct the secretion of an alternative protein by analogous methods.

          The secretory signal sequence contained in the  
25 polypeptides of the present invention can be used to direct other polypeptides into the secretory pathway. The present invention provides for such fusion polypeptides. A signal fusion polypeptide can be made wherein a secretory signal sequence derived from amino acid residues  
30 1-22 of SEQ ID NO:2 is operably linked to another polypeptide using methods known in the art and disclosed herein. The secretory signal sequence contained in the fusion polypeptides of the present invention is preferably fused amino-terminally to an additional peptide to direct  
35 the additional peptide into the secretory pathway. Such constructs have numerous applications known in the art. For example, these novel secretory signal sequence fusion

constructs can direct the secretion of an active component of a normally non-secreted protein, such as a receptor. Such fusions may be used *in vivo* or *in vitro* to direct peptides through the secretory pathway.

5           Cultured mammalian cells are suitable hosts within the present invention. Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection (Wigler et al., Cell 14:725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7:603, 1981; Graham and Van der Eb, Virology 52:456, 10       1973), electroporation (Neumann et al., EMBO J. 1:841-5, 1982), DEAE-dextran mediated transfection (Ausubel et al., ibid.), and liposome-mediated transfection (Hawley-Nelson et al., Focus 15:73, 1993; Ciccarone et al., Focus 15:80, 15       1993, and viral vectors (Miller and Rosman, BioTechniques 7:980-90, 1989; Wang and Finer, Nature Med. 2:714-6, 1996). The production of recombinant polypeptides in cultured mammalian cells is disclosed, for example, by Levinson et al., U.S. Patent No. 4,713,339; Hagen et al., 20       U.S. Patent No. 4,784,950; Palmiter et al., U.S. Patent No. 4,579,821; and Ringold, U.S. Patent No. 4,656,134. Suitable cultured mammalian cells include the COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), BHK (ATCC No. CRL 1632), BHK 570 (ATCC No. CRL 10314), 293 (ATCC No. CRL 25       1573; Graham et al., J. Gen. Virol. 36:59-72, 1977) and Chinese hamster ovary (e.g. CHO-K1; ATCC No. CCL 61) cell lines. Additional suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection, Manassas, VA. In 30       general, strong transcription promoters are preferred, such as promoters from SV-40 or cytomegalovirus. See, e.g., U.S. Patent No. 4,956,288. Other suitable promoters include those from metallothionein genes (U.S. Patent Nos. 4,579,821 and 4,601,978) and the adenovirus major late 35       promoter.

Drug selection is generally used to select for cultured mammalian cells into which foreign DNA has been

inserted. Such cells are commonly referred to as "transfectants". Cells that have been cultured in the presence of the selective agent and are able to pass the gene of interest to their progeny are referred to as

5 "stable transfectants." A preferred selectable marker is a gene encoding resistance to the antibiotic neomycin. Selection is carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems may also be used to increase the expression level of the

10 gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high

15 levels of the products of the introduced genes. A preferred amplifiable selectable marker is dihydrofolate reductase, which confers resistance to methotrexate. Other drug resistance genes (e.g., hygromycin resistance, multi-drug resistance, puromycin acetyltransferase) can

20 also be used. Alternative markers that introduce an altered phenotype, such as green fluorescent protein, or cell surface proteins such as CD4, CD8, Class I MHC, placental alkaline phosphatase may be used to sort transfected cells from untransfected cells by such means

25 as FACS sorting or magnetic bead separation technology.

Other higher eukaryotic cells can also be used as hosts, including plant cells, insect cells and avian cells. The use of *Agrobacterium rhizogenes* as a vector for expressing genes in plant cells has been reviewed by

30 Sinkar et al., J. Biosci. (Bangalore) 11:47-58, 1987. Transformation of insect cells and production of foreign polypeptides therein is disclosed by Guarino et al., U.S. Patent No. 5,162,222 and WIPO publication WO 94/06463. Insect cells can be infected with recombinant baculovirus,

35 commonly derived from *Autographa californica nuclear polyhedrosis virus* (AcNPV). See, King and Possee, The Baculovirus Expression System: A Laboratory Guide,

London, Chapman & Hall; O'Reilly et al., Baculovirus Expression Vectors: A Laboratory Manual, New York, Oxford University Press., 1994; and, Richardson, C. D., Ed., Baculovirus Expression Protocols. Methods in Molecular Biology, Totowa, NJ, Humana Press, 1995. A second method of making recombinant zacr3 baculovirus utilizes a transposon-based system described by Luckow (Luckow et al., J. Virol. 67:4566-79, 1993). This system, which utilizes transfer vectors, is sold in the Bac-to-Bac™ kit (Life Technologies, Rockville, MD). This system utilizes a transfer vector, pFastBac1™ (Life Technologies) containing a Tn7 transposon to move the DNA encoding the zacr3 polypeptide into a baculovirus genome maintained in *E. coli* as a large plasmid called a "bacmid." The pFastBac1™ transfer vector utilizes the AcNPV polyhedrin promoter to drive the expression of the gene of interest, in this case zacr3. However, pFastBac1™ can be modified to a considerable degree. The polyhedrin promoter can be removed and substituted with the baculovirus basic protein promoter (also known as Pcor, p6.9 or MP promoter) which is expressed earlier in the baculovirus infection, and has been shown to be advantageous for expressing secreted proteins. See, Hill-Perkins and Possee, J. Gen. Virol. 71:971-6, 1990; Bonning et al., J. Gen. Virol. 75:1551-6, 1994; and, Chazenbalk, and Rapoport, J. Biol. Chem. 270:1543-9, 1995. In such transfer vector constructs, a short or long version of the basic protein promoter can be used. Moreover, transfer vectors can be constructed which replace the native zacr3 secretory signal sequences with secretory signal sequences derived from insect proteins. For example, a secretory signal sequence from Ecdysteroid Glucosyltransferase (EGT), honey bee Melittin (Invitrogen, Carlsbad, CA), or baculovirus gp67 (PharMingen, San Diego, CA) can be used in constructs to replace the native zacr3 secretory signal sequence. In addition, transfer vectors

can include an in-frame fusion with DNA encoding an epitope tag at the C- or N-terminus of the expressed zacrp3 polypeptide, for example, a Glu-Glu epitope tag (Grussenmeyer et al., Proc. Natl. Acad. Sci. 82:7952-4, 1985). Using a technique known in the art, a transfer vector containing zacrp3 is transformed into *E. coli*, and screened for bacmids which contain an interrupted lacZ gene indicative of recombinant baculovirus. The bacmid DNA containing the recombinant baculovirus genome is isolated, using common techniques, and used to transfect *Spodoptera frugiperda* cells, e.g. Sf9 cells. Recombinant virus that expresses zacrp3 is subsequently produced. Recombinant viral stocks are made by methods commonly used the art.

The recombinant virus is used to infect host cells, typically a cell line derived from the fall armyworm, *Spodoptera frugiperda*. See, in general, Glick and Pasternak, Molecular Biotechnology: Principles and Applications of Recombinant DNA, ASM Press, Washington, D.C., 1994. Another suitable cell line is the High Five<sup>TM</sup> cell line (Invitrogen) derived from *Trichoplusia ni* (U.S. Patent #5,300,435). Commercially available serum-free media are used to grow and maintain the cells. Suitable media are Sf900 II<sup>TM</sup> (Life Technologies) or ESF 921<sup>TM</sup> (Expression Systems) for the Sf9 cells; and Ex-cello405<sup>TM</sup> (JRH Biosciences, Lenexa, KS) or Express FiveO<sup>TM</sup> (Life Technologies) for the *T. ni* cells. The cells are grown up from an inoculation density of approximately  $2-5 \times 10^5$  cells to a density of  $1-2 \times 10^6$  cells at which time a recombinant viral stock is added at a multiplicity of infection (MOI) of 0.1 to 10, more typically near 3. Procedures used are generally described in available laboratory manuals (King and Possee, ibid.; O'Reilly et al., ibid.; Richardson, ibid.). Subsequent purification

of the zacrp3 polypeptide from the supernatant can be achieved using methods described herein.

Fungal cells, including yeast cells, can also be used within the present invention. Yeast species of particular interest in this regard include *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Pichia methanolica*. Methods for transforming *S. cerevisiae* cells with exogenous DNA and producing recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S. Patent No. 4,599,311; Kawasaki et al., U.S. Patent No. 4,931,373; Brake, U.S. Patent No. 4,870,008; Welch et al., U.S. Patent No. 5,037,743; and Murray et al., U.S. Patent No. 4,845,075. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (e.g., leucine). A preferred vector system for use in *Saccharomyces cerevisiae* is the *POT1* vector system disclosed by Kawasaki et al. (U.S. Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Suitable promoters and terminators for use in yeast include those from glycolytic enzyme genes (see, e.g., Kawasaki, U.S. Patent No. 4,599,311; Kingsman et al., U.S. Patent No. 4,615,974; and Bitter, U.S. Patent No. 4,977,092) and alcohol dehydrogenase genes. See also U.S. Patents Nos. 4,990,446; 5,063,154; 5,139,936 and 4,661,454. Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia methanolica*, *Pichia guilliermondii* and *Candida maltosa* are known in the art. See, for example, Gleeson et al., J. Gen. Microbiol. 132:3459-65, 1986 and Cregg, U.S. Patent No. 4,882,279. *Aspergillus* cells may be utilized according to the methods of McKnight et al., U.S. Patent No. 4,935,349. Methods for transforming *Acremonium chrysogenum* are disclosed by Sumino et al., U.S. Patent No. 5,162,228. Methods for

transforming *Neurospora* are disclosed by Lambowitz, U.S. Patent No. 4,486,533.

The use of *Pichia methanolica* as host for the production of recombinant proteins is disclosed in WIPO Publications WO 97/17450, WO 97/17451, WO 98/02536, and WO 98/02565. DNA molecules for use in transforming *P. methanolica* will commonly be prepared as double-stranded, circular plasmids, which are preferably linearized prior to transformation. For polypeptide production in *P. methanolica*, it is preferred that the promoter and terminator in the plasmid be that of a *P. methanolica* gene, such as a *P. methanolica* alcohol utilization gene (*AUG1* or *AUG2*). Other useful promoters include those of the dihydroxyacetone synthase (DHAS), formate dehydrogenase (FMD), and catalase (CAT) genes. To facilitate integration of the DNA into the host chromosome, it is preferred to have the entire expression segment of the plasmid flanked at both ends by host DNA sequences. A preferred selectable marker for use in *Pichia methanolica* is a *P. methanolica* *ADE2* gene, which encodes phosphoribosyl-5-aminoimidazole carboxylase (AIRC; EC 4.1.1.21), which allows *ade2* host cells to grow in the absence of adenine. For large-scale, industrial processes where it is desirable to minimize the use of methanol, it is preferred to use host cells in which both methanol utilization genes (*AUG1* and *AUG2*) are deleted. For production of secreted proteins, host cells deficient in vacuolar protease genes (*PEP4* and *PRB1*) are preferred. Electroporation is used to facilitate the introduction of a plasmid containing DNA encoding a polypeptide of interest into *P. methanolica* cells. It is preferred to transform *P. methanolica* cells by electroporation using an exponentially decaying, pulsed electric field having a field strength of from 2.5 to 4.5 kV/cm, preferably about 3.75 kV/cm, and a time constant ( $\tau$ ) of from 1 to 40 milliseconds, most preferably about 20 milliseconds.

Prokaryotic host cells, including strains of the bacteria *Escherichia coli*, *Bacillus* and other genera are also useful host cells within the present invention. Techniques for transforming these hosts and expressing foreign DNA sequences cloned therein are well known in the art (see, e.g., Sambrook et al., ibid.). When expressing a zacrp3 polypeptide in bacteria such as *E. coli*, the polypeptide may be retained in the cytoplasm, typically as insoluble granules, or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed, and the granules are recovered and denatured using, for example, guanidine isothiocyanate or urea. The denatured polypeptide can then be refolded and dimerized by diluting the denaturant, such as by dialysis against a solution of urea and a combination of reduced and oxidized glutathione, followed by dialysis against a buffered saline solution. In the latter case, the polypeptide can be recovered from the periplasmic space in a soluble and functional form by disrupting the cells (by, for example, sonication or osmotic shock) to release the contents of the periplasmic space and recovering the protein, thereby obviating the need for denaturation and refolding.

Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen host cells. A variety of suitable media, including defined media and complex media, are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals. Media may also contain such components as growth factors or serum, as required. The growth medium will generally select for cells containing the exogenously added DNA by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell.

Expressed recombinant zacrp3 polypeptides (or chimeric zacrp3 polypeptides) can be purified using fractionation and/or conventional purification methods and media. Ammonium sulfate precipitation and acid or chaotrope extraction may be used for fractionation of samples. Exemplary purification steps may include hydroxyapatite, size exclusion, FPLC and reverse-phase high performance liquid chromatography. Suitable chromatographic media include derivatized dextrans, agarose, cellulose, polyacrylamide, specialty silicas, and the like. PEI, DEAE, QAE and Q derivatives are preferred. Exemplary chromatographic media include those media derivatized with phenyl, butyl, or octyl groups, such as Phenyl-Sepharose FF (Pharmacia), Toyopearl butyl 650 (Toso Haas, Montgomeryville, PA), Octyl-Sepharose (Pharmacia) and the like; or polyacrylic resins, such as Amberchrom CG 71 (Toso Haas) and the like. Suitable solid supports include glass beads, silica-based resins, cellulosic resins, agarose beads, cross-linked agarose beads, polystyrene beads, cross-linked polyacrylamide resins and the like that are insoluble under the conditions in which they are to be used. These supports may be modified with reactive groups that allow attachment of proteins by amino groups, carboxyl groups, sulfhydryl groups, hydroxyl groups and/or carbohydrate moieties. Examples of coupling chemistries include cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, hydrazide activation, and carboxyl and amino derivatives for carbodiimide coupling chemistries. These and other solid media are well known and widely used in the art, and are available from commercial suppliers. Methods for binding receptor polypeptides to support media are well known in the art. Selection of a particular method is a matter of routine design and is determined in part by the properties of the chosen support. See, for example, Affinity

Chromatography: Principles & Methods, Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988.

The polypeptides of the present invention can be isolated by exploitation of their structural or binding properties. For example, immobilized metal ion adsorption (IMAC) chromatography can be used to purify histidine-rich proteins or proteins having a His tag. Briefly, a gel is first charged with divalent metal ions to form a chelate (Sulkowski, Trends in Biochem. 3:1-7, 1985). Histidine-rich proteins will be adsorbed to this matrix with differing affinities, depending upon the metal ion used, and will be eluted by competitive elution, lowering the pH, or use of strong chelating agents. Other methods of purification include purification of glycosylated proteins by lectin affinity chromatography and ion exchange chromatography (Methods in Enzymol., Vol. 182, "Guide to Protein Purification", Deutscher, (ed.), Acad. Press, San Diego, 1990, pp. 529-39). Within an additional preferred embodiment of the invention, a fusion of the polypeptide of interest and an affinity tag (e.g., maltose-binding protein, FLAG, Glu-Glu, an immunoglobulin domain) may be constructed to facilitate purification as is discussed in greater detail in the Example sections below.

Protein refolding (and optionally, reoxidation) procedures may be advantageously used. It is preferred to purify the protein to >80% purity, more preferably to >90% purity, even more preferably >95%, and particularly preferred is a pharmaceutically pure state, that is greater than 99.9% pure with respect to contaminating macromolecules, particularly other proteins and nucleic acids, and free of infectious and pyrogenic agents. Preferably, a purified protein is substantially free of other proteins, particularly other proteins of animal origin.

Zacrp3 polypeptides or fragments thereof may also be prepared through chemical synthesis by methods well known in the art. Such zacrp3 polypeptides may be

monomers or multimers; glycosylated or non-glycosylated; pegylated or non-pegylated; and may or may not include an initial methionine amino acid residue.

A ligand-binding polypeptide, such as a zacrp3-binding polypeptide, can also be used for purification of ligand. The polypeptide is immobilized on a solid support, such as beads of agarose, cross-linked agarose, glass, cellulosic resins, silica-based resins, polystyrene, cross-linked polyacrylamide, or like materials that are stable under the conditions of use. Methods for linking polypeptides to solid supports are known in the art, and include amine chemistry, cyanogen bromide activation, N-hydroxysuccinimide activation, epoxide activation, sulfhydryl activation, and hydrazide activation. The resulting medium will generally be configured in the form of a column, and fluids containing ligand are passed through the column one or more times to allow ligand to bind to the ligand-binding polypeptide. The ligand is then eluted using changes in salt concentration, chaotropic agents (guanidine HCl), or pH to disrupt ligand-receptor binding.

An assay system that uses a ligand-binding receptor (or an antibody, one member of a complement/anti-complement pair) or a binding fragment thereof, and a commercially available biosensor instrument (BIAcore™, Pharmacia Biosensor, Piscataway, NJ) may be advantageously employed. Such receptor, antibody, member of a complement/anti-complement pair or fragment is immobilized onto the surface of a receptor chip. Use of this instrument is disclosed by Karlsson, J. Immunol. Methods 145:229-40, 1991 and Cunningham and Wells, J. Mol. Biol. 234:554-63, 1993. A receptor, antibody, member or fragment is covalently attached, using amine or sulfhydryl chemistry, to dextran fibers that are attached to gold film within the flow cell. A test sample is passed through the cell. If a ligand, epitope, or opposite member of the complement/anti-complement pair is present

in the sample, it will bind to the immobilized receptor, antibody or member, respectively, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon resonance of the gold film.

5 This system allows the determination of on- and off-rates, from which binding affinity can be calculated, and assessment of stoichiometry of binding.

Ligand-binding polypeptides can also be used within other assay systems known in the art. Such systems  
10 include Scatchard analysis for determination of binding affinity (see Scatchard, Ann. NY Acad. Sci. 51: 660-72, 1949) and calorimetric assays (Cunningham et al., Science 253:545-48, 1991; Cunningham et al., Science 245:821-25, 1991).

15 The invention also provides anti-zacrp3 antibodies. Antibodies to zacrp3 can be obtained, for example, using as an antigen the product of a zacrp3 expression vector, or zacrp3 isolated from a natural source. Particularly useful anti-zacrp3 antibodies "bind  
20 specifically" with zacrp3. Antibodies are considered to be specifically binding if the antibodies bind to a zacrp3 polypeptide, peptide or epitope with a binding affinity ( $K_a$ ) of  $10^6 \text{ M}^{-1}$  or greater, preferably  $10^7 \text{ M}^{-1}$  or greater, more preferably  $10^8 \text{ M}^{-1}$  or greater, and most preferably  
25  $10^9 \text{ M}^{-1}$  or greater. The binding affinity of an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis (Scatchard, Ann. NY Acad. Sci. 51:660, 1949). Suitable antibodies include antibodies that bind with zacrp3 in particular domains.

30 Anti-zacrp3 antibodies can be produced using antigenic zacrp3 epitope-bearing peptides and polypeptides. Antigenic epitope-bearing peptides and polypeptides of the present invention contain a sequence of at least nine, preferably between 15 to about 30 amino  
35 acids contained within SEQ ID NO:2. However, peptides or polypeptides comprising a larger portion of an amino acid sequence of the invention, containing from 30 to 50 amino

acids, or any length up to and including the entire amino acid sequence of a polypeptide of the invention, also are useful for inducing antibodies that bind with zacrp3. It is desirable that the amino acid sequence of the epitope-bearing peptide is selected to provide substantial solubility in aqueous solvents (i.e., the sequence includes relatively hydrophilic residues, while hydrophobic residues are preferably avoided). Hydrophilic peptides can be predicted by one of skill in the art from a hydrophobicity plot, see for example, Hopp and Woods (Proc. Nat. Acad. Sci. USA 78:3824-8, 1981) and Kyte and Doolittle (J. Mol. Biol. 157: 105-142, 1982). Moreover, amino acid sequences containing proline residues may be also be desirable for antibody production.

Polyclonal antibodies to recombinant zacrp3 protein or to zacrp3 isolated from natural sources can be prepared using methods well-known to those of skill in the art. See, for example, Green et al., "Production of Polyclonal Antisera," in Immunochemical Protocols (Manson, ed.), pages 1-5 (Humana Press 1992), and Williams et al., "Expression of foreign proteins in *E. coli* using plasmid vectors and purification of specific polyclonal antibodies," in DNA Cloning 2: Expression Systems, 2nd Edition, Glover et al. (eds.), page 15 (Oxford University Press 1995). The immunogenicity of a zacrp3 polypeptide can be increased through the use of an adjuvant, such as alum (aluminum hydroxide) or Freund's complete or incomplete adjuvant. Polypeptides useful for immunization also include fusion polypeptides, such as fusions of zacrp3 or a portion thereof with an immunoglobulin polypeptide or with maltose binding protein. The polypeptide immunogen may be a full-length molecule or a portion thereof. If the polypeptide portion is "hapten-like," such portion may be advantageously joined or linked to a macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA) or tetanus toxoid) for immunization.

Although polyclonal antibodies are typically raised in animals such as horses, cows, dogs, chicken, rats, mice, rabbits, hamsters, guinea pigs, goats, or sheep, an anti-zacrp3 antibody of the present invention  
5 may also be derived from a subhuman primate antibody. General techniques for raising diagnostically and therapeutically useful antibodies in baboons may be found, for example, in Goldenberg et al., international patent publication No. WO 91/11465, and in Losman et al., Int. J.  
10 Cancer 46:310, 1990. Antibodies can also be raised in transgenic animals such as transgenic sheep, cows, goats or pigs, and can also be expressed in yeast and fungi in modified forms as well as in mammalian and insect cells.

Alternatively, monoclonal anti-zacrp3 antibodies  
15 can be generated. Rodent monoclonal antibodies to specific antigens may be obtained by methods known to those skilled in the art (see, for example, Kohler et al., Nature 256:495 (1975), Coligan et al. (eds.), Current Protocols in Immunology, Vol. 1, pages 2.5.1-2.6.7 (John  
20 Wiley & Sons 1991), Picksley et al., "Production of monoclonal antibodies against proteins expressed in *E. coli*," in DNA Cloning 2: Expression Systems, 2nd Edition, Glover et al. (eds.), page 93 (Oxford University Press 1995)).

25 Briefly, monoclonal antibodies can be obtained by injecting mice with a composition comprising a zacrp3 gene product, verifying the presence of antibody production by removing a serum sample, removing the spleen to obtain B-lymphocytes, fusing the B-lymphocytes with  
30 myeloma cells to produce hybridomas, cloning the hybridomas, selecting positive clones which produce antibodies to the antigen, culturing the clones that produce antibodies to the antigen, and isolating the antibodies from the hybridoma cultures.

35 In addition, an anti-zacrp3 antibody of the present invention may be derived from a human monoclonal antibody. Human monoclonal antibodies are obtained from

transgenic mice that have been engineered to produce specific human antibodies in response to antigenic challenge. In this technique, elements of the human heavy and light chain locus are introduced into strains of mice  
5 derived from embryonic stem cell lines that contain targeted disruptions of the endogenous heavy chain and light chain loci. The transgenic mice can synthesize human antibodies specific for human antigens, and the mice can be used to produce human antibody-secreting hybridomas.  
10 Methods for obtaining human antibodies from transgenic mice are described, for example, by Green et al., Nature Genet. 7:13, 1994, Lonberg et al., Nature 368:856, 1994, and Taylor et al., Int. Immun. 6:579, 1994.

Monoclonal antibodies can be isolated and  
15 purified from hybridoma cultures by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography (see, for example, Coligan at pages 2.7.1-2.7.12 and pages  
20 2.9.1-2.9.3; Baines et al., "Purification of Immunoglobulin G (IgG)," in Methods in Molecular Biology, Vol. 10, pages 79-104 (The Humana Press, Inc. 1992)).

For particular uses, it may be desirable to prepare fragments of anti-zacrp3 antibodies. Such  
25 antibody fragments can be obtained, for example, by proteolytic hydrolysis of the antibody. Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods. As an illustration, antibody fragments can be produced by  
30 enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted  $F(ab')_2$ . This fragment can be further cleaved using a thiol reducing agent to produce 3.5S Fab' monovalent fragments. Optionally, the cleavage reaction can be performed using a blocking group for the sulfhydryl  
35 groups that result from cleavage of disulfide linkages. As an alternative, an enzymatic cleavage using pepsin produces two monovalent Fab fragments and an Fc fragment

directly. These methods are described, for example, by Goldenberg, U.S. patent No. 4,331,647, Nisonoff et al., Arch Biochem. Biophys. 89:230, 1960, Porter, Biochem. J. 73:119, 1959, Edelman et al., in Methods in Enzymology 5 Vol. 1, page 422 (Academic Press 1967), and by Coligan, ibid.

Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other 10 enzymatic, chemical or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

For example, Fv fragments comprise an association of V<sub>H</sub> and V<sub>L</sub> chains. This association can be 15 noncovalent, as described by Inbar et al., Proc. Natl. Acad. Sci. USA 69:2659, 1972. Alternatively, the variable chains can be linked by an intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde (see, for example, Sandhu, Crit. Rev. Biotech. 12:437, 1992).

20 The Fv fragments may comprise V<sub>H</sub> and V<sub>L</sub> chains which are connected by a peptide linker. These single-chain antigen binding proteins (scFv) are prepared by constructing a structural gene comprising DNA sequences encoding the V<sub>H</sub> and V<sub>L</sub> domains which are connected by an 25 oligonucleotide. The structural gene is inserted into an expression vector which is subsequently introduced into a host cell, such as *E. coli*. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing 30 scFvs are described, for example, by Whitlow et al., Methods: A Companion to Methods in Enzymology 2:97, 1991, also see, Bird et al., Science 242:423, 1988, Ladner et al., U.S. Patent No. 4,946,778, Pack et al., Bio/Technology 11:1271, 1993, and Sandhu, ibid.

35 As an illustration, a scFV can be obtained by exposing lymphocytes to zacrp3 polypeptide *in vitro*, and selecting antibody display libraries in phage or similar

vectors (for instance, through use of immobilized or labeled zacrp3 protein or peptide). Genes encoding polypeptides having potential zacrp3 polypeptide binding domains can be obtained by screening random peptide libraries displayed on phage (phage display) or on bacteria, such as *E. coli*. Nucleotide sequences encoding the polypeptides can be obtained in a number of ways, such as through random mutagenesis and random polynucleotide synthesis. These random peptide display libraries can be used to screen for peptides which interact with a known target which can be a protein or polypeptide, such as a ligand or receptor, a biological or synthetic macromolecule, or organic or inorganic substances. Techniques for creating and screening such random peptide display libraries are known in the art (Ladner et al., U.S. Patent No. 5,223,409, Ladner et al., U.S. Patent No. 4,946,778, Ladner et al., U.S. Patent No. 5,403,484, Ladner et al., U.S. Patent No. 5,571,698, and Kay et al., Phage Display of Peptides and Proteins (Academic Press, Inc. 1996)) and random peptide display libraries and kits for screening such libraries are available commercially, for instance from Clontech (Palo Alto, CA), Invitrogen Inc. (San Diego, CA), New England Biolabs, Inc. (Beverly, MA), and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ). Random peptide display libraries can be screened using the zacrp3 sequences disclosed herein to identify proteins which bind to zacrp3.

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for example, by using the polymerase chain reaction to synthesize the variable region from RNA of antibody-producing cells (see, for example, Larrick et al., Methods: A Companion to Methods in Enzymology 2:106; 1991), Courtenay-Luck, "Genetic Manipulation of Monoclonal

Antibodies," in Monoclonal Antibodies: Production, Engineering and Clinical Application, Ritter et al. (eds.), page 166 (Cambridge University Press, 1995), and Ward et al., "Genetic Manipulation and Expression of  
5 Antibodies," in Monoclonal Antibodies: Principles and Applications, Birch et al., (eds.), page 137 (Wiley-Liss, Inc. 1995)).

Alternatively, an anti-zacrp3 antibody may be derived from a "humanized" monoclonal antibody. Humanized  
10 monoclonal antibodies are produced by transferring mouse complementary determining regions from heavy and light variable chains of the mouse immunoglobulin into a human variable domain. Typical residues of human antibodies are then substituted in the framework regions of the murine  
15 counterparts. The use of antibody components derived from humanized monoclonal antibodies obviates potential problems associated with the immunogenicity of murine constant regions. General techniques for cloning murine immunoglobulin variable domains are described, for  
20 example, by Orlandi et al., Proc. Nat. Acad. Sci. USA 86:3833, 1989. Techniques for producing humanized monoclonal antibodies are described, for example, by Jones et al., Nature 321:522, 1986, Carter et al., Proc. Nat. Acad. Sci. USA 89:4285, 1992, Sandhu, Crit. Rev. Biotech.  
25 12:437, 1992, Singer et al., J. Immun. 150:2844, 1993, Sudhir (ed.), Antibody Engineering Protocols (Humana Press, Inc. 1995), Kelley, "Engineering Therapeutic Antibodies," in Protein Engineering: Principles and Practice, Cleland et al. (eds.), pages 399-434 (John Wiley  
30 & Sons, Inc. 1996), and by Queen et al., U.S. Patent No. 5,693,762 (1997).

Polyclonal anti-idiotypic antibodies can be prepared by immunizing animals with anti-zacrp3 antibodies or antibody fragments, using standard techniques. See,  
35 for example, Green et al., "Production of Polyclonal Antisera," in Methods In Molecular Biology: Immunochemical Protocols, Manson (ed.), pages 1-12 (Humana Press 1992).

Also, see Coligan, ibid. at pages 2.4.1-2.4.7. Alternatively, monoclonal anti-idiotypic antibodies can be prepared using anti-zacrp3 antibodies or antibody fragments as immunogens with the techniques, described  
5 above. As another alternative, humanized anti-idiotypic antibodies or subhuman primate anti-idiotypic antibodies can be prepared using the above-described techniques. Methods for producing anti-idiotypic antibodies are described, for example, by Irie, U.S. Patent No.  
10 5,208,146, Greene, et. al., U.S. Patent No. 5,637,677, and Varthakavi and Minocha, J. Gen. Virol. 77:1875, 1996.

Genes encoding polypeptides having potential zacrp3 polypeptide binding domains, "binding proteins", can be obtained by screening random or directed peptide  
15 libraries displayed on phage (phage display) or on bacteria, such as *E. coli*. Nucleotide sequences encoding the polypeptides can be obtained in a number of ways, such as through random mutagenesis and random polynucleotide synthesis. Alternatively, constrained phage display  
20 libraries can also be produced. These peptide display libraries can be used to screen for peptides which interact with a known target which can be a protein or polypeptide, such as a ligand or receptor, a biological or synthetic macromolecule, or organic or inorganic  
25 substances. Techniques for creating and screening such peptide display libraries are known in the art (Ladner et al., US Patent NO. 5,223,409; Ladner et al., US Patent NO. 4,946,778; Ladner et al., US Patent NO. 5,403,484 and Ladner et al., US Patent NO. 5,571,698) and peptide  
30 display libraries and kits for screening such libraries are available commercially, for instance from Clontech (Palo Alto, CA), Invitrogen Inc. (San Diego, CA), New England Biolabs, Inc. (Beverly, MA) and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ). Peptide display  
35 libraries can be screened using the zacrp3 sequences disclosed herein to identify proteins which bind to zacrp3. These "binding proteins" which interact with

zacrp3 polypeptides can be used essentially like an antibody.

A variety of assays known to those skilled in the art can be utilized to detect antibodies and/or  
5 binding proteins which specifically bind to zacrp3 proteins or peptides. Exemplary assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.), Cold Spring Harbor Laboratory Press, 1988. Representative examples of such assays include: concurrent  
10 immunoelectrophoresis, radioimmunoassay, radioimmuno-precipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assay, inhibition or competition assay, and sandwich assay. In addition, antibodies can be screened for binding to wild-type versus mutant zacrp3  
15 protein or polypeptide.

Antibodies and binding proteins to zacrp3 may be used for tagging cells that express zacrp3; for isolating zacrp3 by affinity purification; for diagnostic assays for determining circulating levels of zacrp3 polypeptides; for  
20 detecting or quantitating soluble zacrp3 as marker of underlying pathology or disease; in analytical methods employing FACS; for screening expression libraries; for generating anti-idiotypic antibodies; and as neutralizing antibodies or as antagonists to block zacrp3 polypeptide  
25 modulation of spermatogenesis or like activity *in vitro* and *in vivo*. Suitable direct tags or labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers, chemiluminescent markers, magnetic particles and the like; indirect tags or labels may  
30 feature use of biotin-avidin or other complement/anti-complement pairs as intermediates. Moreover, antibodies to zacrp3 or fragments thereof may be used *in vitro* to detect denatured zacrp3 or fragments thereof in assays, for example, Western Blots or other assays known in the  
35 art.

Antibodies or polypeptides herein can also be directly or indirectly conjugated to drugs, toxins,

radionuclides and the like, and these conjugates used for in vivo diagnostic or therapeutic applications. For instance, polypeptides or antibodies of the present invention can be used to identify or treat tissues or organs that express a corresponding anti-complementary molecule (receptor or antigen, respectively, for instance). More specifically, zacrp3 polypeptides or anti-zacrp3 antibodies, or bioactive fragments or portions thereof, can be coupled to detectable or cytotoxic molecules and delivered to a mammal having cells, tissues or organs that express the anti-complementary molecule.

An additional aspect of the present invention provides methods for identifying agonists or antagonists of the zacrp3 polypeptides disclosed above, which agonists or antagonists may have valuable properties as discussed further herein. Within one embodiment, there is provided a method of identifying zacrp3 polypeptide agonists, comprising providing cells responsive thereto, culturing the cells in the presence of a test compound and comparing the cellular response with the cell cultured in the presence of the zacrp3 polypeptide, and selecting the test compounds for which the cellular response is of the same type.

Within another embodiment, there is provided a method of identifying antagonists of zacrp3 polypeptide, comprising providing cells responsive to a zacrp3 polypeptide, culturing a first portion of the cells in the presence of zacrp3 polypeptide, culturing a second portion of the cells in the presence of the zacrp3 polypeptide and a test compound, and detecting a decrease in a cellular response of the second portion of the cells as compared to the first portion of the cells. In addition to those assays disclosed herein, samples can be tested for inhibition of zacrp3 activity within a variety of assays designed to measure receptor binding or the stimulation/inhibition of zacrp3-dependent cellular responses. For example, zacrp3-responsive cell lines can

be transfected with a reporter gene construct that is responsive to a zacrp3-stimulated cellular pathway. Reporter gene constructs of this type are known in the art, and will generally comprise a zacrp3-DNA response element operably linked to a gene encoding an assayable protein, such as luciferase. DNA response elements can include, but are not limited to, cyclic AMP response elements (CRE), hormone response elements (HRE), insulin response element (IRE) (Nasrin et al., Proc. Natl. Acad. Sci. USA 87:5273-7, 1990) and serum response elements (SRE) (Shaw et al. Cell 56: 563-72, 1989). Cyclic AMP response elements are reviewed in Roestler et al., J. Biol. Chem. 263 (19):9063-6, 1988 and Habener, Molec. Endocrinol. 4 (8):1087-94, 1990. Hormone response elements are reviewed in Beato, Cell 56:335-44; 1989. Candidate compounds, solutions, mixtures or extracts are tested for the ability to inhibit the activity of zacrp3 on the target cells as evidenced by a decrease in zacrp3 stimulation of reporter gene expression. Assays of this type will detect compounds that directly block zacrp3 binding to cell-surface receptors, as well as compounds that block processes in the cellular pathway subsequent to receptor-ligand binding. In the alternative, compounds or other samples can be tested for direct blocking of zacrp3 binding to receptor using zacrp3 tagged with a detectable label (e.g., <sup>125</sup>I, biotin, horseradish peroxidase, FITC, and the like). Within assays of this type, the ability of a test sample to inhibit the binding of labeled zacrp3 to the receptor is indicative of inhibitory activity, which can be confirmed through secondary assays. Receptors used within binding assays may be cellular receptors or isolated, immobilized receptors.

Based on homology to other adipocyte complement related proteins, zacrp3 polypeptides, fragments, fusions, agonists or antagonists can be used to modulate energy balance in mammals or to protect endothelial cells from injury. With regard to modulating energy balance, zacrp3

polypeptides modulate cellular metabolic reactions. Such metabolic reactions include adipogenesis, gluconeogenesis, glycogenolysis, lipogenesis, glucose uptake, protein synthesis, thermogenesis, oxygen utilization and the like.

5 Zacrp3 polypeptides may also find use as neurotransmitters or as modulators of neurotransmission, as indicated by expression of the polypeptide in tissues associated with the sympathetic or parasympathetic nervous system. In this regard, zacrp3 polypeptides may find utility in

10 modulating nutrient uptake, as demonstrated, for example, by 2-deoxy-glucose uptake in the brain or the like.

Among other methods known in the art or described herein, mammalian energy balance may be evaluated by monitoring one or more of the following

15 metabolic functions: adipogenesis, gluconeogenesis, glycogenolysis, lipogenesis, glucose uptake, protein synthesis, thermogenesis, oxygen utilization or the like. These metabolic functions are monitored by techniques (assays or animal models) known to one of ordinary skill

20 in the art, as is more fully set forth below. For example, the glucoregulatory effects of insulin are predominantly exerted in the liver, skeletal muscle and adipose tissue. Insulin binds to its cellular receptor in these three tissues and initiates tissue-specific actions

25 that result in, for example, the inhibition of glucose production and the stimulation of glucose utilization. In the liver, insulin stimulates glucose uptake and inhibits gluconeogenesis and glycogenolysis. In skeletal muscle and adipose tissue, insulin acts to stimulate the uptake,

30 storage and utilization of glucose.

Art-recognized methods exist for monitoring all of the metabolic functions recited above. Thus, one of ordinary skill in the art is able to evaluate zacrp3 polypeptides, fragments, fusion proteins, antibodies,

35 agonists and antagonists for metabolic modulating functions. Exemplary modulating techniques are set forth below.

Adipogenesis, gluconeogenesis and glycogenolysis are interrelated components of mammalian energy balance, which may be evaluated by known techniques using, for example, *ob/ob* mice or *db/db* mice. The *ob/ob* mice are  
5 inbred mice that are homozygous for an inactivating mutation at the *ob* (obese) locus. Such *ob/ob* mice are hyperphagic and hypometabolic, and are believed to be deficient in production of circulating OB protein. The *db/db* mice are inbred mice that are homozygous for an  
10 inactivating mutation at the *db* (diabetes) locus. The *db/db* mice display a phenotype similar to that of *ob/ob* mice, except *db/db* mice also display a diabetic phenotype. Such *db/db* mice are believed to be resistant to the effects of circulating OB protein. Also, various *in vitro*  
15 methods of assessing these parameters are known in the art.

Insulin-stimulated lipogenesis, for example, may be monitored by measuring the incorporation of  $^{14}\text{C}$ -acetate into triglyceride (Mackall et al. J. Biol. Chem. 251:6462-  
20 4, 1976) or triglyceride accumulation (Kletzien et al., Mol. Pharmacol. 41:393-8, 1992).

Glucose uptake may be evaluated, for example, in an assay for insulin-stimulated glucose transport. Non-transfected, differentiated L6 myotubes (maintained in the  
25 absence of G418) are placed in DMEM containing 1 g/l glucose, 0.5 or 1.0% BSA, 20 mM Hepes, and 2 mM glutamine. After two to five hours of culture, the medium is replaced with fresh, glucose-free DMEM containing 0.5 or 1.0% BSA, 20 mM Hepes, 1 mM pyruvate, and 2 mM glutamine.  
30 Appropriate concentrations of insulin or IGF-1, or a dilution series of the test substance, are added, and the cells are incubated for 20-30 minutes.  $^3\text{H}$  or  $^{14}\text{C}$ -labeled deoxyglucose is added to  $\approx 50$   $\mu\text{M}$  final concentration, and the cells are incubated for approximately 10-30 minutes.  
35 The cells are then quickly rinsed with cold buffer (e.g. PBS), then lysed with a suitable lysing agent (e.g. 1% SDS or 1 N NaOH). The cell lysate is then evaluated by

counting in a scintillation counter. Cell-associated radioactivity is taken as a measure of glucose transport after subtracting non-specific binding as determined by incubating cells in the presence of cytocholasin b, an inhibitor of glucose transport. Other methods include those described by, for example, Manchester et al., Am. J. Physiol. 266 (Endocrinol. Metab. 29):E326-E333, 1994 (insulin-stimulated glucose transport).

Protein synthesis may be evaluated, for example, by comparing precipitation of <sup>35</sup>S-methionine-labeled proteins following incubation of the test cells with <sup>35</sup>S-methionine and <sup>35</sup>S-methionine and a putative modulator of protein synthesis.

Thermogenesis may be evaluated as described by B. Stanley in The Biology of Neuropeptide Y and Related Peptides, W. Colmers and C. Wahlestedt (eds.), Humana Press, Ottawa, 1993, pp. 457-509; C. Billington et al., Am. J. Physiol. 260:R321, 1991; N. Zarjevski et al., Endocrinology 133:1753, 1993; C. Billington et al., Am. J. Physiol. 266:R1765, 1994; Heller et al., Am. J. Physiol. 252(4 Pt 2): R661-7, 1987; and Heller et al., Am. J. Physiol. 245: R321-8, 1983. Also, metabolic rate, which may be measured by a variety of techniques, is an indirect measurement of thermogenesis.

Oxygen utilization may be evaluated as described by Heller et al., Pflugers Arch 369: 55-9, 1977. This method also involved an analysis of hypothalamic temperature and metabolic heat production. Oxygen utilization and thermoregulation have also been evaluated in humans as described by Haskell et al., J. Appl. Physiol. 51: 948-54, 1981.

Neurotransmission functions may be evaluated by monitoring 2-deoxy-glucose uptake in the brain. This parameter is monitored by techniques (assays or animal models) known to one of ordinary skill in the art, for example, autoradiography. Useful monitoring techniques are described, for example, by Kilduff et al., J.

Neurosci. 10 2463-75, 1990, with related techniques used to evaluate the "hibernating heart" as described in Gerber et al. Circulation 94: 651-8, 1996, and Fallavollita et al., Circulation 95: 1900-9, 1997.

5           In addition, zacrp3 polypeptides, fragments, fusions agonists or antagonists thereof may be therapeutically useful for anti-microbial applications. For example, complement component C1q plays a role in host defense against infectious agents, such as bacteria and  
10           viruses. C1q is known to exhibit several specialized functions. For example, C1q triggers the complement cascade via interaction with bound antibody or C-reactive protein (CRP). Also, C1q interacts directly with certain bacteria, RNA viruses, mycoplasma, uric acid crystals, the  
15           lipid A component of bacterial endotoxin and membranes of certain intracellular organelles. C1q binding to the C1q receptor is believed to promote phagocytosis. C1q also appears to enhance the antibody formation aspect of the host defense system. See, for example, Johnston, Pediatr.  
20           Infect. Dis. J. 12(11): 933-41, 1993. Thus, soluble C1q-like molecules may be useful as anti-microbial agents, promoting lysis or phagocytosis of infectious agents.

          Zacrp3 fragments as well as zacrp3 polypeptides, fusion proteins, agonists, antagonists or antibodies may  
25           be evaluated with respect to their anti-microbial properties according to procedures known in the art. See, for example, Barsum et al., Eur. Respir. J. 8(5): 709-14, 1995; Sandovsky-Losica et al., J. Med. Vet. Mycol (England) 28(4): 279-87, 1990; Mehentee et al., J. Gen.  
30           Microbiol. (England) 135 (Pt. 8): 2181-8, 1989; Segal and Savage, J. Med. Vet. Mycol. 24: 477-9, 1986 and the like. If desired, the performance of zacrp3 in this regard can be compared to proteins known to be functional in this regard, such as proline-rich proteins, lysozyme,  
35           histatins, lactoperoxidase or the like. In addition, zacrp3 fragments, polypeptides, fusion proteins, agonists, antagonists or antibodies may be evaluated in combination

with one or more anti-microbial agents to identify synergistic effects. One of ordinary skill in the art will recognize that the anti-microbial properties of zacrp3 polypeptides, fragments, fusion proteins, agonists, antagonists and antibodies may be similarly evaluated.

As neurotransmitters or neurotransmission modulators, zacrp3 polypeptide fragments as well as zacrp3 polypeptides, fusion proteins, agonists, antagonists or antibodies of the present invention may also modulate calcium ion concentration, muscle contraction, hormone secretion, DNA synthesis or cell growth, inositol phosphate turnover, arachidonate release, phospholipase-C activation, gastric emptying, human neutrophil activation or ADCC capability, superoxide anion production and the like. Evaluation of these properties can be conducted by known methods, such as those set forth herein.

The impact of zacrp3 polypeptide, fragment, fusion, antibody, agonist or antagonist on intracellular calcium level may be assessed by methods known in the art, such as those described by Dobrzanski et al., Regulatory Peptides 45: 341-52, 1993, and the like. The impact of zacrp3 polypeptide, fragment, fusion, agonist or antagonist on muscle contraction may be assessed by methods known in the art, such as those described by Smits & Lebeuvre, J. Auton. Pharmacol. 14: 383-92, 1994, Belloli et al., J. Vet. Pharmacol. Therap. 17: 379-83, 1994, Maggi et al., Regulatory Peptides 53: 259-74, 1994, and the like. The impact of zacrp3 polypeptide, fragment, fusion, agonist or antagonist on hormone secretion may be assessed by methods known in the art, such as those for prolactin release described by Henriksen et al., J. Recep. Sig. Transd. Res. 15(1-4): 529-41, 1995, and the like. The impact of zacrp3 polypeptide, fragment, fusion, agonist or antagonist on DNA synthesis or cell growth may be assessed by methods known in the art, such as those described by Dobrzanski et al., Regulatory Peptides 45: 341-52, 1993, and the like. The impact of zacrp3 polypeptide, fragment,

fusion, agonist or antagonist on inositol phosphate turnover may be assessed by methods known in the art, such as those described by Dobrzanski et al., Regulatory Peptides 45: 341-52, 1993, and the like.

5 Also, the impact of zacrp3 polypeptide, fragment, fusion, agonist or antagonist on arachidonate release may be assessed by methods known in the art, such as those described by Dobrzanski et al., Regulatory Peptides 45: 341-52, 1993, and the like. The impact of  
10 zacrp3 polypeptide, fragment, fusion, agonist or antagonist on phospholipase-C activation may be assessed by methods known in the art, such as those described by Dobrzanski et al., Regulatory Peptides 45: 341-52, 1993, and the like. The impact of zacrp3 polypeptide, fragment,  
15 fusion, agonist or antagonist on gastric emptying may be assessed by methods known in the art, such as those described by Varga et al., Eur. J. Pharmacol. 286: 109-112, 1995, and the like. The impact of zacrp3 polypeptide, fragment, fusion, agonist or antagonist on  
20 human neutrophil activation and ADCC capability may be assessed by methods known in the art, such as those described by Wozniak et al., Immunology 78: 629-34, 1993, and the like. The impact of zacrp3 polypeptide, fragment, fusion, agonist or antagonist on superoxide anion  
25 production may be assessed by methods known in the art, such as those described by Wozniak et al., Immunology 78: 629-34, 1993, and the like.

Collagen is a potent inducer of platelet aggregation. This poses risks to patients recovering from  
30 vascular injuries. Inhibitors of collagen-induced platelet aggregation would be useful for blocking the binding of platelets to collagen-coated surfaces and reducing associated collagen-induced platelet aggregation. Clq is a component of the complement pathway and has been found  
35 to stimulate defense mechanisms as well as trigger the generation of toxic oxygen species that can cause tissue damage (Tenner, Behring Inst. Mitt. 93:241-53, 1993). Clq

binding sites are found on platelets. Clq, independent of an immune binding partner, has been found to inhibit platelet aggregation but not platelet adhesion or shape change. The amino terminal region of Clq shares homology  
5 with collagen (Peerschke and Ghebrehiwet, J. Immunol. 145:2984-88, 1990). Inhibition of Clq and the complement pathway can be determined using methods disclosed herein or know in the art, such as described in Suba and Csako, J. Immunol. 117:304-9, 1976.

10 The impact of zacrp3 polypeptide, fragments, fusions, agonists or antagonists on collagen-mediated platelet adhesion, activation and aggregation may be evaluated using methods described herein or known in the art, such as the platelet aggregation assay (Chiang et al., Thrombosis Res. 37:605-12, 1985) and platelet  
15 adhesion assays (Peerschke and Ghebrehiwet, J. Immunol. 144:221-25, 1990). Assays for platelet adhesion to collagen and inhibition of collagen-induced platelet aggregation can be measured using methods described in  
20 Keller et al., J. Biol. Chem. 268:5450-6, 1993; Waxman and Connolly, J. Biol. Chem. 268:5445-9, 1993; Noeske-Jungblut et al., J. Biol. Chem. 269:5050-3 or 1994 Deckmyn et al., Blood 85:712-9, 1995.

The impact of zacrp3 polypeptide, fragments, fusions, agonists or antagonists on vasodilation of aortic rings can be measured according to the methods of Dainty et al., J. Pharmacol. 100:767, 1990 and Rhee et al., Neurotox. 16:179, 1995.

Various in vitro and in vivo models are  
30 available for assessing the effects of zacrp3 polypeptides, fragments, fusion proteins, antibodies, agonists and antagonists on ischemia and reperfusion injury. See for example, Shandelya et al., Circulation 88:2812-26, 1993; Weisman et al., Science 249:146-151, 1991; Buerke et al., Circulation 91:393-402, 1995;  
35 Horstick et al., Circulation 95:701-8, 1997 and Burke et al., J. Phar. Exp. Therp. 286:429-38, 1998. An ex vivo

hamster platelet aggregation assay is described by Deckmyn et al., ibid. Bleeding times in hamsters and baboons can be measured following injection of zacrp3 polypeptides using the model described by Deckmyn et al., ibid. The  
5 formation of thrombus in response to administration of proteins of the present invention can be measured using the hamster femoral vein thrombosis model is provided by Deckmyn et al., ibid. Changes in platelet adhesion under flow conditions following administration of zacrp3 can be  
10 measured using the method described in Harsfalvi et al., Blood 85:705-11, 1995.

Complement inhibition and wound healing can be zacrp3 polypeptides, fragments, fusion proteins, antibodies, agonists or antagonists be assayed alone or in  
15 combination with other know-inhibitors of collagen-induced platelet activation and aggregation, such as palldipin, moubatin or calin, for example.

Zacrp3 polypeptides, fragments, fusion proteins, antibodies, agonists or antagonists can be evaluated using  
20 methods described herein or known in the art, such as healing of dermal layers in pigs (Lynch et al., Proc. Natl. Acad. Sci. USA 84: 7696-700, 1987) and full-thickness skin wounds in genetically diabetic mice (Greenhalgh et al., Am. J. Pathol. 136: 1235-46, 1990),  
25 for example. The polypeptides of the present invention can be assayed alone or in combination with other known complement inhibitors as described above.

Radiation hybrid mapping is a somatic cell genetic technique developed for constructing high-  
30 resolution, contiguous maps of mammalian chromosomes (Cox et al., Science 250:245-50, 1990). Partial or full knowledge of a gene's sequence allows the designing of PCR primers suitable for use with chromosomal radiation hybrid mapping panels. Commercially available radiation hybrid  
35 mapping panels which cover the entire human genome, such as the Stanford G3 RH Panel and the GeneBridge 4 RH Panel (Research Genetics, Inc., Huntsville, AL), are available.

These panels enable rapid, PCR based, chromosomal localizations and ordering of genes, sequence-tagged sites (STSS), and other nonpolymorphic- and polymorphic markers within a region of interest. This includes establishing  
5 directly proportional physical distances between newly discovered genes of interest and previously mapped markers. The precise knowledge of a gene's position can be useful in a number of ways including: 1) determining if a sequence is part of an existing contig and obtaining  
10 additional surrounding genetic sequences in various forms such as YAC-, BAC- or cDNA clones, 2) providing a possible candidate gene for an inheritable disease which shows linkage to the same chromosomal region, and 3) for cross-referencing model organisms such as mouse which may  
15 be beneficial in helping to determine what function a particular gene might have.

The results showed linkage of Zacrp3 to the human chromosome 5 framework marker SHGC-56588 with a LOD score of 15.58 and at a distance of 0 cR\_10000 from the  
20 marker. The use of surrounding markers positions Zacrp3 in the 5p12 region on the integrated LDB human chromosome 5 map. The present invention also provides reagents which will find use in diagnostic applications. For example, the zacrp3 gene, a probe comprising zacrp3 DNA or RNA, or  
25 a subsequence thereof can be used to determine if the zacrp3 gene is present on chromosome 5 or if a mutation has occurred. Detectable chromosomal aberrations at the zacrp3 gene locus include, but are not limited to, aneuploidy, gene copy number changes, insertions,  
30 deletions, restriction site changes and rearrangements. These aberrations can occur within the coding sequence, within introns, or within flanking sequences, including upstream promoter and regulatory regions, and may be manifested as physical alterations within a coding  
35 sequence or changes in gene expression level.

In general, these diagnostic methods comprise the steps of (a) obtaining a genetic sample from a

patient; (b) incubating the genetic sample with a polynucleotide probe or primer as disclosed above, under conditions wherein the polynucleotide will hybridize to complementary polynucleotide sequence, to produce a first reaction product; and (iii) comparing the first reaction product to a control reaction product. A difference between the first reaction product and the control reaction product is indicative of a genetic abnormality in the patient. Genetic samples for use within the present invention include genomic DNA, cDNA, and RNA. The polynucleotide probe or primer can be RNA or DNA, and will comprise a portion of SEQ ID NO:1, the complement of SEQ ID NO:1, or an RNA equivalent thereof. Suitable assay methods in this regard include molecular genetic techniques known to those in the art, such as restriction fragment length polymorphism (RFLP) analysis, short tandem repeat (STR) analysis employing PCR techniques, ligation chain reaction (Barany, PCR Methods and Applications 1:5-16, 1991), ribonuclease protection assays, and other genetic linkage analysis techniques known in the art (Sambrook et al., ibid.; Ausubel et. al., ibid.; Marian, Chest 108:255-65, 1995). Ribonuclease protection assays (see, e.g., Ausubel et al., ibid., ch. 4) comprise the hybridization of an RNA probe to a patient RNA sample, after which the reaction product (RNA-RNA hybrid) is exposed to RNase. Hybridized regions of the RNA are protected from digestion. Within PCR assays, a patient's genetic sample is incubated with a pair of polynucleotide primers, and the region between the primers is amplified and recovered. Changes in size or amount of recovered product are indicative of mutations in the patient. Another PCR-based technique that can be employed is single strand conformational polymorphism (SSCP) analysis (Hayashi, PCR Methods and Applications 1:34-8, 1991).

Zacrp3 polypeptides may be used in the analysis of energy efficiency of a mammal. Zacrp3 polypeptides found in serum or tissue samples may be indicative of a

mammals ability to store food, with more highly efficient mammals tending toward obesity. More specifically, the present invention contemplates methods for detecting zacrp3 polypeptide comprising:

5 exposing a sample possibly containing zacrp3 polypeptide to an antibody attached to a solid support, wherein said antibody binds to an epitope of a zacrp3 polypeptide;

washing said immobilized antibody-polypeptide to  
10 remove unbound contaminants;

exposing the immobilized antibody-polypeptide to a second antibody directed to a second epitope of a zacrp3 polypeptide, wherein the second antibody is associated with a detectable label; and

15 detecting the detectable label. The concentration of zacrp3 polypeptide in the test sample appears to be indicative of the energy efficiency of a mammal. This information can aid nutritional analysis of a mammal. Potentially, this information may be useful in  
20 identifying and/or targeting energy deficient tissue.

A further aspect of the invention provides a method for studying insulin. Such methods of the present invention comprise incubating adipocytes in a culture medium comprising zacrp3 polypeptide, monoclonal antibody,  
25 agonist or antagonist thereof  $\pm$  insulin and observing changes in adipocyte protein secretion or differentiation.

Anti-microbial protective agents may be directly acting or indirectly acting. Such agents operating via membrane association or pore forming mechanisms of action  
30 directly attach to the offending microbe. Anti-microbial agents can also act via an enzymatic mechanism, breaking down microbial protective substances or the cell wall/membrane thereof. Anti-microbial agents, capable of inhibiting microorganism proliferation or action or of  
35 disrupting microorganism integrity by either mechanism set forth above, are useful in methods for preventing contamination in cell culture by microbes susceptible to

that anti-microbial activity. Such techniques involve culturing cells in the presence of an effective amount of said zacrp3 polypeptide or an agonist or antagonist thereof.

5           Also, zacrp3 polypeptides or agonists thereof may be used as cell culture reagents in *in vitro* studies of exogenous microorganism infection, such as bacterial, viral or fungal infection. Such moieties may also be used in *in vivo* animal models of infection.

10           The present invention also provides methods of studying mammalian cellular metabolism. Such methods of the present invention comprise incubating cells to be studied, for example, human vascular endothelial cells,  $\pm$  zacrp3 polypeptide, monoclonal antibody, agonist or  
15 antagonist thereof and observing changes in adipogenesis, gluconeogenesis, glycogenolysis, lipogenesis, glucose uptake, or the like.

          An additional aspect of the invention provides a method for studying dimerization or oligomerization. Such  
20 methods of the present invention comprise incubating zacrp3 polypeptides or fragments or fusion proteins thereof containing a collagen-like domain alone or in combination with other polypeptides bearing collagen-like domains and observing the associations formed between the  
25 collagen like domains. Such associations are indicated by HPLC, circular dichroism or the like.

          Zacrp3 polypeptides, fragments, fusion proteins, antibodies, agonists or antagonists of the present invention can be used in methods for promoting blood flow  
30 within the vasculature of a mammal by reducing the number of platelets that adhere and are activated and the size of platelet aggregates. Used to such an end, Zacrp3 can be administered prior to, during or following an acute vascular injury in the mammal. Vascular injury may be due  
35 to vascular reconstruction, including but not limited to, angioplasty, coronary artery bypass graft, microvascular repair or anastomosis of a vascular graft. Also

contemplated are vascular injuries due to trauma, stroke or aneurysm. In other preferred methods the vascular injury is due to plaque rupture, degradation of the vasculature, complications associated with diabetes and atherosclerosis. Plaque rupture in the coronary artery induces heart attack and in the cerebral artery induces stroke. Use of zacrp3 polypeptides, fragments, fusion proteins, antibodies, agonists or antagonists in such methods would also be useful for ameliorating whole system diseases of the vasculature associated with the immune system, such as disseminated intravascular coagulation (DIC) and SIDs. Additionally the complement inhibiting activity would be useful for treating non-vasculature immune diseases such as arteriolosclerosis.

A correlation has been found between the presence of C1q in localized ischemic myocardium and the accumulation of leukocytes following coronary occlusion and reperfusion. Release of cellular components following tissue damage triggers complement activation which results in toxic oxygen products that may be the primary cause of myocardial damage (Rossen et al., Circ. Res. 62:572-84, 1998 and Tenner, ibid.). Blocking the complement pathway was found to protect ischemic myocardium from reperfusion injury (Buerke et al., J. Pharm. Exp. Therp. 286:429-38, 1998). Proteins having complement inhibition and C1q binding activity would be useful for such purposes.

Collagen and C1q binding capabilities of adipocyte complement related protein homologs such as zacrp3 would be useful to pacify damaged collagenous tissues preventing platelet adhesion, activation or aggregation, and the activation of inflammatory processes which lead to the release of toxic oxygen products. By rendering the exposed tissue inert towards such processes as complement activity, thrombotic activity and immune activation, reduces the injurious effects of ischemia and reperfusion. In particular, such injuries would include trauma injury ischemia, intestinal strangulation, and

injury associated with pre- and post-establishment of blood flow. Such polypeptides would be useful in the treatment of cardiopulmonary bypass ischemia and revascularization, myocardial infarction and post trauma  
5 vasospasm, such as stroke or percutaneous transluminal angioplasty as well as accidental or surgical-induced vascular trauma.

Additionally such collagen- and C1q-binding polypeptides would be useful to pacify prosthetic  
10 biomaterials and surgical equipment to render the surface of the materials inert towards complement activation, thrombotic activity or immune activation. Such materials include, but are not limited to, collagen or collagen fragment-coated biomaterials, gelatin-coated biomaterials,  
15 fibrin-coated biomaterials, fibronectin-coated biomaterials, heparin-coated biomaterials, collagen and gel-coated stents, arterial grafts, synthetic heart valves, artificial organs or any prosthetic application exposed to blood that will bind zsig37 at greater than  $1 \times 10^8$ .  
20 Coating such materials can be done using methods known in the art, see for example, Rubens, US Patent No. 5,272,074.

Complement and C1q play a role in inflammation. The complement activation is initiated by binding of C1q  
25 to immunoglobulins (Johnston, Pediatr. Infect. Dis. J. 12:933-41, 1993; Ward and Ghetie, Therap. Immunol. 2:77-94, 1995). Inhibitors of C1q and complement would be useful as anti-inflammatory agents. Such application can be made to prevent infection. Additionally, such  
30 inhibitors can be administered to an individual suffering from inflammation mediated by complement activation and binding of immune complexes to C1q. Inhibitors of C1q and complement would be useful in methods of mediating wound repair, enhancing progression in wound healing by  
35 overcoming impaired wound healing. Progression in wound healing would include, for example, such elements as a

reduction in inflammation, fibroblasts recruitment, wound retraction and reduction in infection.

Ability of tumor cells to bind to collagen may contribute to the metastasis of tumors. Inhibitors of collagen binding are also useful for mediating the adhesive interactions and metastatic spread of tumors (Noeske-Jungbult et al., US Patent No. 5,723,312).

In addition, zacrp3 polypeptides, fragments, fusions agonists or antagonists thereof may be therapeutically useful for anti-microbial applications. For example, complement component C1q plays a role in host defense against infectious agents, such as bacteria and viruses. C1q is known to exhibit several specialized functions. For example, C1q triggers the complement cascade via interaction with bound antibody or C-reactive protein (CRP). Also, C1q interacts directly with certain bacteria, RNA viruses, mycoplasma, uric acid crystals, the lipid A component of bacterial endotoxin and membranes of certain intracellular organelles. C1q binding to the C1q receptor is believed to promote phagocytosis. C1q also appears to enhance the antibody formation aspect of the host defense system. See, for example, Johnston, Pediatr. Infect. Dis. J. 12(11): 933-41, 1993. Thus, soluble C1q-like molecules may be useful as anti-microbial agents, promoting lysis or phagocytosis of infectious agents.

The positively charged, extracellular, triple helix, collagenous domains of C1q and macrophage scavenger receptor were determined to play a role in ligand binding and were shown to have a broad binding specificity for polyanions (Acton et al., J. Biol. Chem. 268:3530-37, 1993). Lysophospholipid growth factor (lysophosphatidic acid, LPA) and other mitogenic anions localize at the site of damaged tissues and assist in wound repair. LPA exerts many biological effects including activation of platelets and up-regulation of matrix assembly. It is thought that LPA synergizes with other blood coagulation factors and mediates wound healing.

The collagenous domains of proteins such as C1q and macrophage scavenger receptor are known to bind acidic phospholipids such as LPA. The interaction of zacrp3 polypeptides, fragments, fusions, agonists or antagonists with mitogenic anions such as LPA can be determined using assays known in the art, see for example, Acton et al., ibid. Inhibition of inflammatory processes by polypeptides and antibodies of the present invention would also be useful in preventing infection at the wound site.

For pharmaceutical use, the proteins of the present invention can be formulated with pharmaceutically acceptable carriers for parenteral, oral, nasal, rectal, topical, transdermal administration or the like, according to conventional methods. In a preferred embodiment administration is made at or near the site of vascular injury. In general, pharmaceutical formulations will include a zacrp3 protein in combination with a pharmaceutically acceptable vehicle, such as saline, buffered saline, 5% dextrose in water or the like. Formulations may further include one or more excipients, preservatives, solubilizers, buffering agents, albumin to prevent protein loss on vial surfaces, etc. Methods of formulation are well known in the art and are disclosed, for example, in Remington: The Science and Practice of Pharmacy, Gennaro, ed., Mack Publishing Co., Easton PA, 19<sup>th</sup> ed., 1995. Therapeutic doses will generally be determined by the clinician according to accepted standards, taking into account the nature and severity of the condition to be treated, patient traits, etc. Determination of dose is within the level of ordinary skill in the art.

As used herein a "pharmaceutically effective amount" of a zsig37 polypeptide, fragment, fusion protein, agonist or antagonist is an amount sufficient to induce a desired biological result. The result can be alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For

example, an effective amount of a zacrp3 polypeptide is that which provides either subjective relief of symptoms or an objectively identifiable improvement as noted by the clinician or other qualified observer. Such an effective amount of a zacrp3 polypeptide would provide, for example, inhibition of collagen-activated platelet activation and the complement pathway, including C1q, increase localized blood flow within the vasculature of a patient and/or reduction in injurious effects of ischemia and reperfusion. Effective amounts of the zacrp3 polypeptides can vary widely depending on the disease or symptom to be treated. The amount of the polypeptide to be administered and its concentration in the formulations, depends upon the vehicle selected, route of administration, the potency of the particular polypeptide, the clinical condition of the patient, the side effects and the stability of the compound in the formulation. Thus, the clinician will employ the appropriate preparation containing the appropriate concentration in the formulation, as well as the amount of formulation administered, depending upon clinical experience with the patient in question or with similar patients. Such amounts will depend, in part, on the particular condition to be treated, age, weight, and general health of the patient, and other factors evident to those skilled in the art. Typically a dose will be in the range of 0.01-100 mg/kg of subject. In applications such as balloon catheters the typical dose range would be 0.05-5 mg/kg of subject. Doses for specific compounds may be determined from in vitro or ex vivo studies in combination with studies on experimental animals. Concentrations of compounds found to be effective in vitro or ex vivo provide guidance for animal studies, wherein doses are calculated to provide similar concentrations at the site of action.

Polynucleotides encoding zacrp3 polypeptides are useful within gene therapy applications where it is desired to increase or inhibit zacrp3 activity. If a

mammal has a mutated or absent zacrp3 gene, the zacrp3 gene can be introduced into the cells of the mammal. In one embodiment, a gene encoding a zacrp3 polypeptide is introduced *in vivo* in a viral vector. Such vectors  
5 include an attenuated or defective DNA virus, such as, but not limited to, herpes simplex virus (HSV), papillomavirus, Epstein Barr virus (EBV), adenovirus, adeno-associated virus (AAV), and the like. Defective viruses, which entirely or almost entirely lack viral  
10 genes, are preferred. A defective virus is not infective after introduction into a cell. Use of defective viral vectors allows for administration to cells in a specific, localized area, without concern that the vector can infect other cells. Examples of particular vectors include, but  
15 are not limited to, a defective herpes simplex virus 1 (HSV1) vector (Kaplitt et al., Molec. Cell. Neurosci. 2:320-30, 1991); an attenuated adenovirus vector, such as the vector described by Stratford-Perricaudet et al., J. Clin. Invest. 90:626-30, 1992; and a defective adeno-associated virus vector (Samulski et al., J. Virol. 61:3096-101, 1987; Samulski et al., J. Virol. 63:3822-8, 1989).

In another embodiment, a zacrp3 gene can be introduced in a retroviral vector, e.g., as described in  
25 Anderson et al., U.S. Patent No. 5,399,346; Mann et al. Cell 33:153, 1983; Temin et al., U.S. Patent No. 4,650,764; Temin et al., U.S. Patent No. 4,980,289; Markowitz et al., J. Virol. 62:1120, 1988; Temin et al., U.S. Patent No. 5,124,263; WIPO Publication WO 95/07358;  
30 and Kuo et al., Blood 82:845, 1993. Alternatively, the vector can be introduced by lipofection *in vivo* using liposomes. Synthetic cationic lipids can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker (Felgner et al., Proc. Natl. Acad. Sci.  
35 USA 84:7413-7, 1987; Mackey et al., Proc. Natl. Acad. Sci. USA 85:8027-31, 1988). The use of lipofection to introduce exogenous genes into specific organs *in vivo* has certain

practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. More particularly, directing transfection to particular cells represents one area of benefit. For instance, directing  
5 transfection to particular cell types would be particularly advantageous in a tissue with cellular heterogeneity, such as the pancreas, liver, kidney, and brain. Lipids may be chemically coupled to other molecules for the purpose of targeting. Targeted peptides  
10 (e.g., hormones or neurotransmitters), proteins such as antibodies, or non-peptide molecules can be coupled to liposomes chemically.

It is possible to remove the target cells from the body; to introduce the vector as a naked DNA plasmid;  
15 and then to re-implant the transformed cells into the body. Naked DNA vectors for gene therapy can be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran,  
20 calcium phosphate precipitation, use of a gene gun or use of a DNA vector transporter. See, e.g., Wu et al., J. Biol. Chem. 267:963-7, 1992; Wu et al., J. Biol. Chem. 263:14621-4, 1988.

Antisense methodology can be used to inhibit  
25 zacrp3 gene transcription, such as to inhibit cell proliferation in vivo. Polynucleotides that are complementary to a segment of a zacrp3-encoding polynucleotide (e.g., a polynucleotide as set forth in SEQ ID NO:1) are designed to bind to zacrp3-encoding mRNA and  
30 to inhibit translation of such mRNA. Such antisense polynucleotides are used to inhibit expression of zacrp3 polypeptide-encoding genes in cell culture or in a subject.

Transgenic mice, engineered to express the  
35 zacrp3 gene, and mice that exhibit a complete absence of zacrp3 gene function, referred to as "knockout mice" (Snouwaert et al., Science 257:1083, 1992), may also be

generated (Lowell et al., Nature 366:740-42, 1993). These mice may be employed to study the zacrp3 gene and the protein encoded thereby in an *in vivo* system.

The invention is further illustrated by the following non-limiting examples.

### EXAMPLES

#### Example 1

#### Chromosomal Assignment and Placement of Zacrp3

10

Zacrp3 was mapped to human chromosome 5 using the commercially available version of the Stanford G3 Radiation Hybrid Mapping Panel (Research Genetics, Inc., Huntsville, AL). The Stanford G3 RH Panel contains PCRable DNAs from each of 83 radiation hybrid clones of the whole human genome, plus two control DNAs (the RM donor and the A3 recipient). A publicly available WWW server (<http://shgc-www.stanford.edu>) allows chromosomal localization of markers.

20

For the mapping of zacrp3 with the Stanford G3 RH Panel, 20  $\mu$ l reactions were set up in a 96-well microtiter plate (Stratagene, La Jolla, CA) and used in a RoboCycler Gradient 96 thermal cycler (Stratagene). Each of the 85 PCR reactions consisted of 2  $\mu$ l 10X KlenTaq PCR reaction buffer (Clontech Laboratories, Inc., Palo Alto, CA), 1.6  $\mu$ l dNTPs mix (2.5 mM each, PERKIN-ELMER, Foster City, CA), 1  $\mu$ l sense primer, ZC 21,913 (SEQ ID NO:13), 1  $\mu$ l antisense primer, ZC 21,914 (SEQ ID NO:14), 2  $\mu$ l RediLoad (Research Genetics, Inc.), 0.4  $\mu$ l 50X Advantage KlenTaq Polymerase Mix (Clontech Laboratories, Inc.), 25 ng of DNA from an individual hybrid clone or control and ddH<sub>2</sub>O for a total volume of 20  $\mu$ l. The reactions were overlaid with an equal amount of mineral oil and sealed. The PCR cycler conditions were as follows: an initial 1 cycle 5 minute denaturation at 94°C, 35 cycles of a 45 seconds denaturation at 94°C, 45 seconds annealing at 62°C and 1 minute and 15 seconds extension at 72°C, followed by

35

a final 1 cycle extension of 7 minutes at 72°C. The reactions were separated by electrophoresis on a 2% agarose gel (Life Technologies, Gaithersburg, MD).

The results showed linkage of Zacrp3 to the human chromosome 5 framework marker SHGC-56588 with a LOD score of 15.58 and at a distance of 0 cR\_10000 from the marker. The use of surrounding markers positions Zacrp3 in the 5p12 region on the integrated LDB human chromosome 5 map (The Genetic Location Database, University of Southhampton, WWW server: [http://cedar.genetics.soton.ac.uk/public\\_html/](http://cedar.genetics.soton.ac.uk/public_html/)).

#### Example 2 Baculovirus Expression of Zacrp3

15

An expression vector, pzacrp3cee, was prepared to express human zacrp3 polypeptides having a carboxy-terminal Glu-Glu tag, in insect cells.

#### 20 A. Construction of pzacrp3cee

A 766 bp fragment containing sequence for zacrp3 (SEQ ID NO:1) and a polynucleotide sequence encoding BamHI and XbaI restriction sites on the 5' and 3' ends, respectively, was generated by PCR amplification from a plasmid containing zacrp3 cDNA using primers ZC23377 (SEQ ID NO:15) and ZC23378 (SEQ ID NO:16). The PCR reaction conditions were as follows: 1 cycle of 94°C for 4 minutes, followed by 25 cycles of 94°C for 45 seconds; 50°C for 45 seconds, and 72°C for 2 minutes; 1 cycle at 72°C for 10 min; followed by a 4°C soak. The fragment was visualized by gel electrophoresis (1% Seaplaque/1% NuSieve). The band was excised, diluted to 0.5% agarose with 2 mM MgCl<sub>2</sub>, melted at 65°C and ligated into an BamHI/XbaI digested baculovirus expression donor vector, pZBV32L. The pZBV32L vector is a modification of the pFastBac1™ (Life Technologies) expression vector, where the polyhedron

promoter has been removed and replaced with the late activating Basic Protein Promoter and the coding sequence for the Glu-Glu tag (SEQ ID NO:17) as well as a stop signal is inserted at the 3' end of the multiple cloning region). About 11 nanograms of the restriction digested zacrp3 insert and about 23 ng of the corresponding vector were ligated overnight at 16°C. The ligation mix was diluted 3 fold in TE (10 mM Tris-HCl, pH 7.5 and 1 mM EDTA) and 4 fmol of the diluted ligation mix was transformed into DH5 $\alpha$  Library Efficiency competent cells (Life Technologies) according to manufacturer's direction by heat shock for 45 seconds in a 42°C waterbath. The transformed DNA and cells were diluted in 450  $\mu$ l of SOC media (2% Bacto Tryptone, 0.5% Bacto Yeast Extract, 10 ml 1M NaCl, 1.5 mM KCl, 10 mM MgCl<sub>2</sub>, 10 mM MgSO<sub>4</sub> and 20 mM glucose) and plated onto LB plates containing 100  $\mu$ g/ml ampicillin. Clones were analyzed by restriction digests and 1  $\mu$ l of the positive clone was transformed into 20  $\mu$ l DH10Bac Max Efficiency competent cells (GIBCO-BRL, Gaithersburg, MD) according to manufacturer's instruction, by heat shock for 45 seconds in a 42°C waterbath. The transformed cells were then diluted in 980  $\mu$ l SOC media (2% Bacto Tryptone, 0.5% Bacto Yeast Extract, 10 ml 1M NaCl, 1.5 mM KCl, 10 mM MgCl<sub>2</sub>, 10 mM MgSO<sub>4</sub> and 20 mM glucose) out grown in shaking incubator at 37°C for four hours and plated onto Luria Agar plates containing 50  $\mu$ g/ml kanamycin, 7  $\mu$ g/ml gentamicin, 10  $\mu$ g/ml tetracycline, IPTG and Blue Gal. The plated cells were incubated for 48 hours at 37°C. A color selection was used to identify those cells having zacrp3cee encoding donor insert that had incorporated into the plasmid (referred to as a "bacmid"). Those colonies, which were white in color, were picked for analysis. Bacmid DNA was isolated from positive colonies using the QiaVac Miniprep8 system (Qiagen) according the manufacturer's directions. Clones

were screened for the correct insert by amplifying DNA using primers to the transposable element in the bacmid via PCR using primers ZC447 (SEQ ID NO:18) and ZC976 (SEQ ID NO:19). The PCR reaction conditions were as follows:  
5 35 cycles of 94°C for 45 seconds, 50°C for 45 seconds, and 72°C for 5 minutes; 1 cycle at 72°C for 10 min.; followed by 4°C soak. The PCR product was run on a 1% agarose gel to check the insert size. Those having the correct insert were used to transfect *Spodoptera frugiperda* (Sf9) cells.

10

#### B. Transfection

Sf9 cells were seeded at  $5 \times 10^6$  cells per 35 mm plate and allowed to attach for 1 hour at 27°C. Five microliters of bacmid DNA was diluted with 100 µl Sf-900  
15 II SFM (Life Technologies). Six µl of CellFECTIN Reagent (Life Technologies) was diluted with 100 µl Sf-900 II SFM. The bacmid DNA and lipid solutions were gently mixed and incubated 30-45 minutes at room temperature. The media from one plate of cells were aspirated, the cells were  
20 washed 1X with 2 ml fresh Sf-900 II SFM media. Eight hundred microliters of Sf-900 II SFM was added to the lipid-DNA mixture. The wash media was aspirated and the DNA-lipid mix added to the cells. The cells were incubated at 27°C for 4-5 hours. The DNA-lipid mix was  
25 aspirated and 2 ml of Sf-900 II media was added to each plate. The plates were incubated at 27°C, 90% humidity, for 96 hours after which the virus was harvested.

#### C. Primary Amplification

30 Sf9 cells were grown in 50 ml Sf-900 II SFM in a 125 ml shake flask to an approximate density of  $0.41-0.52 \times 10^5$  cells/ml. They were then infected with 150 µl of the virus stock from above and incubated at 27°C for 3 days after which time the virus was harvested according to  
35 standard methods known in the art.

Example 3  
Purification of Baculovirus Expressed Glu-Glu-tagged  
zacrp3 polypeptides

5

Unless otherwise noted, all operations were carried out at 4°C. A mixture of protease inhibitors were added to a 2 liter sample of conditioned media from C-terminal Glu-Glu (EE) tagged zacrp3 baculovirus-infected Sf9 cells to final concentrations of 2.5 mM ethylenediaminetetraacetic acid (EDTA, Sigma Chemical Co. St. Louis, MO), 0.001 mM leupeptin (Boehringer-Mannheim, Indianapolis, IN), 0.001 mM pepstatin (Boehringer-Mannheim) and 0.4 mM Pefabloc (Boehringer-Mannheim). The sample was centrifuged at 10,000 rpm for 30 min at 4°C in a Beckman JLA-10.5 rotor (Beckman Instruments) in a Beckman Avanti J25I centrifuge (Beckman Instruments) to remove cell debris. To the supernatant fraction was added a 50.0 ml sample of anti-EE Sepharose, prepared as described below, and the mixture was gently agitated on a Wheaton (Millville, NJ) roller culture apparatus for 18.0 h at 4°C.

The mixture was poured into a 5.0 x 20.0 cm Econo-Column (Bio-Rad Laboratories) and the gel was washed with 30 column volumes of phosphate buffered saline (PBS). The unretained flow-through fraction was discarded. Once the absorbance of the effluent at 280 nm was less than 0.05, flow through the column was reduced to zero and the anti-EE Sepharose gel was washed with 2.0 column volumes of PBS containing 0.2 mg/ml of EE peptide (AnaSpec, San Jose, CA). The peptide used has the sequence Glu-Tyr-Met-Pro-Val-Asp (SEQ ID NO:20). After 1.0 hour at 4°C, flow was resumed and the eluted protein was collected. This fraction was referred to as the peptide elution. The anti-EE Sepharose gel was washed with 2.0 column volumes of 0.1 M glycine, pH 2.5, and the glycine wash was collected separately. The pH of the

glycine-eluted fraction was adjusted to 7.0 by the addition of a small volume of 10X PBS and stored at 4°C.

The peptide elution was concentrated to 5.0 ml using a 5,000 molecular weight cutoff membrane concentrator (Millipore) according to the manufacturer's instructions. The concentrated peptide elution was separated from free peptide by chromatography on a 1.5 x 50 cm Sephadex G-50 (Pharmacia) column equilibrated in PBS at a flow rate of 1.0 ml/min using a BioCad Sprint HPLC (PerSeptive BioSystems). Two ml fractions were collected and the absorbance at 280 nM was monitored. The first peak of material absorbing at 280 nM and eluting near the void volume of the column was collected. This material represented purified zacrp3CEE and was composed of two major bands of apparent molecular weights.

#### Preparation of anti-EE Sepharose

A 100 ml bed volume of protein G-Sepharose (Pharmacia) was washed 3 times with 100 ml of PBS containing 0.02% sodium azide using a 500 ml Nalgene 0.45 micron filter unit. The gel was washed with 6.0 volumes of 200 mM triethanolamine, pH 8.2 (TEA, Sigma), and an equal volume of EE antibody solution containing 900 mg of antibody was added. After an overnight incubation at 4°C, unbound antibody was removed by washing the resin with 5 volumes of 200 mM TEA as described above. The resin was resuspended in 2 volumes of TEA, transferred to a suitable container, and dimethylpimilimidate-2 HCl (Pierce), dissolved in TEA, was added to a final concentration of 36 mg/ml of gel. The gel was rocked at room temperature for 45 min and the liquid was removed using the filter unit as described above. Nonspecific sites on the gel were then blocked by incubating for 10 minutes at room temperature with 5 volumes of 20 mM ethanolamine in 200 mM TEA. The gel was then washed with 5 volumes of PBS containing 0.02% sodium azide and stored in this solution at 4°C.

Example 4  
Adhesion Molecule Assays

5           Upon stimulation with inflammatory cytokines such as TNF (tumor necrosis factor), human microvascular bone marrow cells (TRBMEC) express cell surface adhesion molecules, including E-selectin (endothelial leukocyte  
10 molecule); and I-CAM (intercellular adhesion molecule).

          The effect of zacrp3 on expression of cell surface adhesion molecules was determined using microvascular bone marrow cells (TRBMEC) in a cell based ELISA according to Ouchi et al., (Circulation 100:2473-7,  
15 1999). Briefly, TRBMEC cells were grown in 96 well, flat bottom plates (Costar, Pleasanton, CA) until confluent. Both wild type control media and baculovirus-expressed zacrp3 media was concentrated 10X before testing (Centricon Centrifugal Filtration Unit 5,000K cutoff,  
20 Millipore Corp., Bedford, MA) according to the manufacturer's instructions. To each well 90 µl of zacrp3-containing media or control media was added, and the plates were incubated at 37°C, 5% CO<sub>2</sub> overnight. The next day, half of the samples received 10 µl of TNFα (10  
25 ng/ml, R&D Systems, Minneapolis, MN), the other samples were untreated, measuring basal expression. The plates were then incubated at 37°C, 5% CO<sub>2</sub> for 4 hours.

          Following incubation, the media was removed from the plates and 50 µl anti-human VCAM antibody (1:1000  
30 dilution of a 1 mg/ml stock, R&D Systems), 50 µl of anti-human ICAM-1 monoclonal antibody (1:1000 dilution of a 1 mg/ml stock, R&D Systems), or 50 µl of anti-human E-selectin antibodies (1:1000 dilution of a 1 mg/ml stock, R&D Systems) were then added to triplicate wells and the  
35 plates were incubated at 37°C, 5% CO<sub>2</sub> for 1 hour.

          The antibody solution was removed and the plates were washed three times in warm RPMI + 5% FBS. Following

the last wash, 100  $\mu$ l/well of an 0.05% gluteraldehyde solution (1:1000 of 50% gluteraldehyde in PBS) was added to the wells and the plates were incubated at room temperature for 10 minutes. The plates were washed three  
5 times with PBS and 50  $\mu$ l/well of secondary antibody (1:1000 dilution of goat  $\alpha$ -mouse IgG whole molecule HRP conjugate, (Sigma Chemical Co., St. Louis, Mo.) was added to all wells. The plates were incubated for one hour at 37°C.

10 The plates were then washed five times with washing buffer (PBS + 0.05% Tween 20) and 100  $\mu$ l/well TMB solution (100  $\mu$ l of 4 mg/ml Tetra methyl benzidine (Sigma) in DMSO, in 10 ml 60 mM Na Acetate pH 5.0 and 100  $\mu$ l 1.2% H<sub>2</sub>O<sub>2</sub>) was added to each well. The plates were allowed to  
15 develop at room temperature for 15-20 minutes at which time the reaction was quenched by adding 100  $\mu$ l/well 1M H<sub>2</sub>SO<sub>4</sub>. Plates were read at 450 nm with reference wavelength of 655 nm.

Zacrp3 showed no effect on ICAM-1 expression.  
20 Zacrp3 did show an effect on VCAM-1 expression. When compared to the maximal TNF response, zacrp3 treated cells showed about 50% inhibition. Zacrp3 also had an effect, although less, 10% inhibition of E-selection expression.

VCAM-1 expression was measured following direct  
25 adenovirus infection of TRBMEC cells. Briefly, TRBMEC cells were directly infected with an adenovirus containing zacrp3 or the parental adenovirus strain. The virus was added at various multiplicities of infection (moi 500, 1,000 and 5,000). Cells were incubated at 37°C, 5% CO<sub>2</sub> for  
30 43 hours. Following infection, the adenovirus-infected cells were challenged with TNF $\alpha$  (1 ng/ml) for four hours. VCAM expression was measured as described above. Inhibition of VCAM-1 expression was about 13% at moi 5000, about 5% at moi 1000 and no effect was seen at moi 500.

35 Adenovirus conditioned media was concentrated 10X (Centricon Centrifugal Filtration Unit 5,000K cutoff, Millipore Corp., Bedford, MA) according to the

manufacturer's instructions followed by heat inactivation at 56°C for 30 minutes). The concentrated heat inactivated samples were assayed as described above. For VCAM-1 and E-selectin, inhibition was 100%. Similar  
5 results were observed when either IL-1 or LPS was used to induce adhesion molecule expression. For ICAM-1, inhibition was reduced to nearly baseline. The experiment was repeated with varying concentrations of zacrp3 heat inactivated adenovirus conditioned media. Complete  
10 inhibition was seen at 5X, 50% inhibition at 2.5X and no inhibition at 0.5X.

A THP-1 monocyte adherence assay according to Ouchi et al., (ibid.) and Cybulsky and Gimbrone, (Science 251:788-91, 1991) showed the same results as were seen for  
15 VCAM-1 above.

## CLAIMS

What is claimed is:

1. An isolated polypeptide comprising a sequence of amino acid residues that is at least 75% identical in amino acid sequence to residues 51-246 of SEQ ID NO:2, wherein said sequence comprises:

Gly-Xaa-Xaa or Gly-Xaa-Pro repeats forming a collagen domain, wherein Xaa is any amino acid; and

a carboxyl-terminal Clq domain comprising 10 beta strands.

2. An isolated polypeptide according to claim 1, wherein said polypeptide that is at least 90% identical in amino acid sequence to residues 23-246 of SEQ ID NO:2.

3. An isolated polypeptide according to claim 1, wherein said collagen domain consists of 15 Gly-Xaa-Xaa repeats and 6 Gly-Xaa-Pro repeats.

4. An isolated polypeptide according to claim 1, wherein said carboxyl-terminal Clq domain comprises the sequence of SEQ ID NO:10.

5. An isolated polypeptide according to claim 1, wherein said carboxy-terminal Clq domain comprises amino acid residues 119-123, 140-142, 148-151, 155-158, 161-173, 175-182, 190-197, 200-212, 217-222 and 236-241 of SEQ ID NO:2.

6. An isolated polypeptide according to claim 1, wherein any differences between said polypeptide and SEQ ID NO:2 are due to conservative amino acid substitutions.

7. An isolated polypeptide according to claim 1, wherein said polypeptide specifically binds with an antibody that specifically binds with a polypeptide consisting of the amino acid sequence of SEQ ID NO:2.

8. An isolated polypeptide according to claim 2, wherein said polypeptide comprises residues 23-246 of SEQ ID NO:2.

9. An isolated polypeptide according to claim 1, wherein said collagen domain consists of amino acid residues 51-113 of SEQ ID NO:2.

10. An isolated polypeptide according to claim 1, wherein said carboxy-terminal Clq domain consists of amino acid residues 114-246 of SEQ ID NO:2.

11. An isolated polypeptide according to claim 1, covalently linked at the amino or carboxyl terminus to a moiety selected from the group consisting of affinity tags, toxins, radionucleotides, enzymes and fluorophores.

12. An isolated polypeptide selected from the group consisting of:

a) a polypeptide consisting of a sequence of amino acid residues that is 80% identical in amino acid sequence to amino acid residue 51 to amino acid residue 113 of SEQ ID NO:2, said polypeptide consisting of Gly-Xaa-Xaa and Gly-Xaa-Pro repeats forming a collagen domain;

b) a polypeptide consisting of a sequence of amino acid residues that is 80% identical in amino acid sequence to amino acid residue 114 to amino acid residue 246 of SEQ ID NO:2 comprising the sequence of SEQ ID NO:5; and

c) a polypeptide consisting of a sequence of amino acid residues that is 80% identical in amino acid sequence to amino acid residue 51 to 246 of SEQ ID NO:2, said polypeptide consisting of Gly-Xaa-Xaa and Gly-Xaa-Pro repeats forming a collagen domain and comprising the sequence of SEQ ID NO:5.

13. A fusion protein comprises a first portion and a second portion joined by a peptide bond, said first portion

consisting of a polypeptide selected from the group consisting of:

a) a polypeptide comprising a sequence of amino acid residues that is at least 75% identical in amino acid sequence to amino acid residue 51 to amino acid residue 246 of SEQ ID NO:2;

b) a polypeptide comprising a sequence of amino acid residues as shown in SEQ ID NO:2 from amino acid residue 1 to amino acid residue 246;

c) a portion of the zacrp3 polypeptide of SEQ ID NO:2, comprising the collagen-like domain or a portion of the collagen-like domain capable of dimerization or oligomerization;

d) a portion of the zacrp3 polypeptide of SEQ ID NO:2, comprising the Clq domain or an active portion of the Clq domain; or

e) a portion of the zacrp3 polypeptide of SEQ ID NO:2 comprising of the collagen-like domain and the Clq domain; and

said second portion comprising another polypeptide.

14. A fusion protein according to claim 13, wherein said first portion is selected from the group consisting of:

a) a polypeptide consisting of the sequence of amino acid residue 51 to amino acid residue 113 of SEQ ID NO:2;

b) a polypeptide consisting of the sequence of amino acid residue 114 to amino acid residue 246 of SEQ ID NO:2;

c) a polypeptide consisting of the sequence of amino acid residue 51 to 246 of SEQ ID NO:2.

15. A fusion protein according to claim 13, wherein said second portion comprises a collagen domain or a Clq domain from an adipocyte complement related protein.

16. A polypeptide according to Claim 1; in combination with a pharmaceutically acceptable vehicle.

17. An antibody or antibody fragment that specifically binds to a polypeptide according to claim 1.

18. An antibody according to claim 17, wherein said antibody is selected from the group consisting of:

- a) polyclonal antibody;
- b) murine monoclonal antibody;
- c) humanized antibody derived from b); and
- d) human monoclonal antibody.

19. An antibody fragment according to claim 17, wherein said antibody fragment is selected from the group consisting of F(ab'), F(ab), Fab', Fab, Fv, scFv, and minimal recognition unit.

20. An anti-idiotypic antibody that specifically binds to said antibody of claim 17.

21. An isolated polynucleotide encoding a polypeptide comprising a sequence of amino acid residues that is at least 75% identical in amino acid sequence to residues 51-246 of SEQ ID NO:2, wherein said sequence comprises:

Gly-Xaa-Xaa or Gly-Xaa-Pro repeats forming a collagen domain, wherein Xaa is any amino acid; and  
a carboxyl-terminal C1q domain comprising 10 beta strands.

22. An isolated polynucleotide according to claim 21, wherein said polypeptide is at least 90% identical in amino acid sequence to residues 23-246 of SEQ ID NO:2.

23. An isolated polynucleotide according to claim 21, wherein said collagen domain consists of 15 Gly-Xaa-Xaa repeats and 6 Gly-Xaa-Pro repeats.

24. An isolated polynucleotide according to claim 21, wherein said carboxyl-terminal Clq domain comprises the sequence of SEQ ID NO:5.

25. An isolated polynucleotide according to claim 21, wherein said carboxy-terminal Clq domain comprises amino acid residues 119-123, 140-142, 148-151, 155-158, 161-173, 175-182, 190-197, 200-212, 217-222 and 236-241 of SEQ ID NO:2.

26. An isolated polynucleotide according to claim 21, wherein any differences between said polypeptide and SEQ ID NO:2 are due to conservative amino acid substitutions.

27. An isolated polynucleotide according to claim 21, wherein said polypeptide specifically binds with an antibody that specifically binds with a polypeptide consisting of the amino acid sequence of SEQ ID NO:2.

28. An isolated polynucleotide according to claim 21, wherein said collagen domain comprises amino acid residues 51-113 of SEQ ID NO:2.

29. An isolated polynucleotide according to claim 21, wherein said carboxy-terminal Clq domain consists of amino acid residues 114-246 of SEQ ID NO:2.

30. An isolated polynucleotide according to claim 21, wherein said polypeptide comprises residues 23-246 of SEQ ID NO:2.

31. An isolated polynucleotide selected from the group consisting of,

a) a sequence of nucleotides from nucleotide 1 to nucleotide 1696 of SEQ ID NO:1;

- b) a sequence of nucleotides from nucleotide 69 to nucleotide 806 of SEQ ID NO:1;
- c) a sequence of nucleotides from nucleotide 135 to nucleotide 806 of SEQ ID NO:1;
- d) a sequence of nucleotides from nucleotide 219 to nucleotide 806 of SEQ ID NO:1;
- e) a sequence of nucleotides from nucleotide 408 to nucleotide 806 of SEQ ID NO:1;
- f) a sequence of nucleotides from nucleotide 69 to nucleotide 407 of SEQ ID NO:1;
- g) a sequence of nucleotides from nucleotide 135 to nucleotide 407 of SEQ ID NO:1;
- h) a sequence of nucleotides from nucleotide 219 to nucleotide 407 of SEQ ID NO:1;
- i) a polynucleotide encoding a polypeptide, said polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 51 to 113 of SEQ ID NO:2;
- j) a polynucleotide encoding a polypeptide, said polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 114 to 246 of SEQ ID NO:2;
- k) a polynucleotide encoding a polypeptide, said polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 51 to 246 of SEQ ID NO:2;
- l) a polynucleotide encoding a polypeptide consisting of a sequence of amino acid residues that is at least 75% identical to a polypeptide consisting of the amino acid sequence of residues 23 to 113 of SEQ ID NO:2;
- m) a polynucleotide that remains hybridized following stringent wash conditions to a polynucleotide consisting of the nucleotide sequence of SEQ ID NO:1, or the complement of SEQ ID NO:1;
- n) nucleotide sequences complementary to a), b), c), d), e), f), g), h), i), j), k), l) or m) and

o) degenerate nucleotide sequences of i), j), k) or l).

32. An isolated polynucleotide encoding a fusion protein comprising a first portion and a second portion joined by a peptide bond, said first portion is selected from the group consisting of:

a) a polypeptide comprising a sequence of amino acid residues that is at least 75% identical in amino acid sequence to amino acid residues 51 to 246 of SEQ ID NO:2;

b) a polypeptide comprising the sequence of amino acid residues 1 to 246 of SEQ ID NO:2;

c) a polypeptide comprising the sequence of amino acid residues 23 to 246 of SEQ ID NO:2;

d) a polypeptide comprising the sequence of amino acid residues 23 to 113 of SEQ ID NO:2;

e) a polypeptide comprising the sequence of amino acid residues 1 to 113 of SEQ ID NO:2;

f) a portion of a polypeptide of SEQ ID NO:2 comprising the collagen-like domain or a portion of the collagen-like domain capable of dimerization or oligomerization;

g) a portion of the polypeptide of SEQ ID NO:2 containing the Clq domain; or

h) a portion of the polypeptide of SEQ ID NO:2 including the collagen-like domain and the Clq domain; and  
said second portion comprising another polypeptide.

33. An isolated polynucleotide consisting of the sequence of nucleotide 1 to nucleotide 738 of SEQ ID NO:10.

34. An expression vector comprising the following operably linked elements:

a transcription promoter;

a DNA segment encoding a polypeptide according to claim 1; and

a transcription terminator.

35. An expression vector according to claim 34, wherein said DNA segment encodes a polypeptide that is at least 90% identical in amino acid sequence to residues 23-246 of SEQ ID NO:2.

36. An expression vector according to claim 34, wherein said collagen domain consists of 15 Gly-Xaa-Xaa repeats and 6 Gly-Xaa-Pro repeats.

37. An expression vector according to claim 34, wherein said carboxyl-terminal Clq domain comprises the sequence of SEQ ID NO:10.

38. An expression vector according to claim 34, wherein said carboxy-terminal Clq domain comprises amino acid residues 119-123, 140-142, 148-151, 155-158, 161-173, 175-182, 190-197, 200-212, 217-222 and 236-241 of SEQ ID NO:2.

39. An expression vector according to claim 34, wherein any differences between said polypeptide and SEQ ID NO:2 are due to conservative amino acid substitutions.

40. An expression vector according to claim 34, wherein said polypeptide specifically binds with an antibody that specifically binds with a polypeptide consisting of the amino acid sequence of SEQ ID NO:2.

41. An expression vector according to claim 34, wherein said collagen domain consists of amino acid residues 51-113 of SEQ ID NO:2.

42. An expression vector according to claim 34, wherein said Clq domain consists of amino acid residues 114-246 of SEQ ID NO:2.

43. An expression vector according to claim 34, wherein said DNA segment encodes a polypeptide comprising residues 23-246 of SEQ ID NO:2.

44. An expression vector according to claim 34; wherein said DNA segment encodes a polypeptide covalently linked at the amino or carboxyl terminus to an affinity tag.

45. An expression vector according to claim 34 wherein said DNA segment further encodes a secretory signal sequence operably linked to said polypeptide.

46. An expression vector according the claim 34, wherein said secretory signal sequence comprises residues 1-22 of SEQ ID NO:2.

47. A cultured cell into which has been introduced an expression vector according to claim 34, wherein said cell expresses said polypeptide encoded by said DNA segment.

48. A method of producing a polypeptide comprising:  
culturing a cell into which has been introduced an expression vector according to claim 34;

whereby said cell expresses said polypeptide encoded by said DNA segment; and

recovering said expressed polypeptide.

1/1

	10	20	30	40	50		
C1QC_HUMAN	MDVGPSSLP	HLGLKLLLLLLLL	LALRGOANT	-----		; 30	
ACR3_HUMAN	MLLLGAVLLLL	LALPGHDQETTT	QGPGVLLPL	PKGACTGWMA	-----	; 41	
HUZACRP3	MLWRQLIYWQ	LLALFFLPFCL	CQDEYMESPQ	TGGLPPDCSK	CCHGDYSFR	; 50	
	60	70	80	90	100		
C1QC_HUMAN	-----G	CYGIPGMPGL	PAPGKDG	YDGLPGPK	GEPGIPAIP	GIRGP ; 71	
ACR3_HUMAN	GIPGHPGH	NGAPGRDGR	DGTPGEK	GEKGDPL	IGPKGDI	GETGVPGAEGP ; 91	
HUZACRP3	GYQGP	PPGPPGIP	PGNHG	NNGNGAT	GHEGAKGE	KGDKGDLGPRGERGQ ; 100	
	110	120	130	140	150		
C1QC_HUMAN	KGQKGE	PGLPGHPG	KNGPMGP	PGVP	PGMGP	IPGEPGEEGRY----- ; 114	
ACR3_HUMAN	RGFPGI	QGRKGEP	-----			; 104	
HUZACRP3	HGPKGE	KGYPGIP	-----			; 113	
	160	170	180	190	200		
C1QC_HUMAN	----KQK	FQSVFTV	TRQTHQP	PAPNSL	IRFNAVL	TNPQGDYDTSTGKFTC ; 160	
ACR3_HUMAN	--GEG	AYVYR	SAFSV	GLET	YVTIPN	MPIRFTKIFYNQNH	DYGSTGKFHC ; 152
HUZACRP3	-----	PELQIA	FMASL	ATHFSN	QNSGII	FSSVETNIGNFFD	VMTGRFGA ; 157
	210	220	230	240	250		
C1QC_HUMAN	KVPGLYY	FVYHAS-	HTANLC	VLLYRSG	---VKV	VTCGHTSKTNQVN--S ; 204	
ACR3_HUMAN	NIPGLYY	FAYHIT	VYMKD	VKVS	LFFKK---	DKAMLFTYDQYQENNV	DQAS ; 198
HUZACRP3	PVSGVY	FFTF	SMMKH	EDVEE	VYVYLM	HNGNTVFS	MYSYEMKGKSD--TSS ; 205
	260	270	280	290	300		
C1QC_HUMAN	GGVLLRL	QVGEEV	WLAVND	YYDM	VGIQG---	SDSVFSGFLLFPD----- ; 245	
ACR3_HUMAN	GSVLLH	LEVGDQ	VWLQVY	GEGER	NGLYAD-	NDNDSTFTGFLLYH	DTN--- ; 244
HUZACRP3	NHAVL	KLAKG	DEVW	LRMG-	NGALH	G	GDHQR---FSTFAGFLLFETK--- ; 246

Fig. 1

## SEQUENCE LISTING

&lt;110&gt; ZymoGenetics, Inc.

&lt;120&gt; ADIPOCYTE-SPECIFIC PROTEIN HOMOLOG ZACRP3

&lt;130&gt; 99-09

&lt;150&gt; 60/130.199

&lt;151&gt; 1999-04-20

&lt;160&gt; 20

&lt;170&gt; FastSEQ for Windows Version 3.0

&lt;210&gt; 1

&lt;211&gt; 1696

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (69)...(806)

&lt;400&gt; 1

```

cccaggaga ccacgctcct ggagctctgc tgtcttctca gggagactct gaggctctgt      60
tgagaatc atg ctt tgg agg cag ctc atc tat tgg caa ctg ctg gct ttg      110
      Met Leu Trp Arg Gln Leu Ile Tyr Trp Gln Leu Leu Ala Leu
            1             5             10

ttt ttc ctc cct ttt tgc ctg tgt caa gat gaa tac atg gag tct cca      158
Phe Phe Leu Pro Phe Cys Leu Cys Gln Asp Glu Tyr Met Glu Ser Pro
      15             20             25             30

caa acc gga gga cta ccc cca gac tgc agt aag tgt tgt cat gga gac      206
Gln Thr Gly Gly Leu Pro Pro Asp Cys Ser Lys Cys Cys His Gly Asp
            35             40             45

tac agc ttt cga ggc tac caa ggc ccc cct ggg cca ccg ggc cct cct      254
Tyr Ser Phe Arg Gly Tyr Gln Gly Pro Pro Gly Pro Pro Gly Pro Pro
            50             55             60

```

ggc att cca gga aac cat gga aac aat ggc aac aat gga gcc act ggt Gly Ile Pro Gly Asn His Gly Asn Asn Gly Asn Asn Gly Ala Thr Gly 65 70 75	302
cat gaa gga gcc aaa ggt gag aag ggc gac aaa ggt gac ctg ggg cct His Glu Gly Ala Lys Gly Glu Lys Gly Asp Lys Gly Asp Leu Gly Pro 80 85 90	350
cga ggg gag cgg ggg cag cat ggc ccc aaa gga gag aag ggc tac ccg Arg Gly Glu Arg Gly Gln His Gly Pro Lys Gly Glu Lys Gly Tyr Pro 95 100 105 110	398
ggg att cca cca gaa ctt cag att gca ttc atg gct tct ctg gca acc Gly Ile Pro Pro Glu Leu Gln Ile Ala Phe Met Ala Ser Leu Ala Thr 115 120 125	446
cac ttc agc aat cag aac agt ggg att atc ttc agc agt gtt gag acc His Phe Ser Asn Gln Asn Ser Gly Ile Ile Phe Ser Ser Val Glu Thr 130 135 140	494
aac att gga aac ttc ttt gat gtc atg act ggt aga ttt ggg gcc cca Asn Ile Gly Asn Phe Phe Asp Val Met Thr Gly Arg Phe Gly Ala Pro 145 150 155	542
gta tca ggt gtg tat ttc ttc acc ttc agc atg atg aag cat gag gat Val Ser Gly Val Tyr Phe Phe Thr Phe Ser Met Met Lys His Glu Asp 160 165 170	590
gtt gag gaa gtg tat gtg tac ctt atg cac aat ggc aac aca gtc ttc Val Glu Glu Val Tyr Val Tyr Leu Met His Asn Gly Asn Thr Val Phe 175 180 185 190	638
agc atg tac agc tat gaa atg aag ggc aaa tca gat aca tcc agc aat Ser Met Tyr Ser Tyr Glu Met Lys Gly Lys Ser Asp Thr Ser Ser Asn 195 200 205	686
cat gct gtg ctg aag cta gcc aaa ggg gat gag gtt tgg ctg cga atg His Ala Val Leu Lys Leu Ala Lys Gly Asp Glu Val Trp Leu Arg Met 210 215 220	734
ggc aat ggc gct ctc cat ggg gac cac caa cgc ttc tcc acc ttt gca Gly Asn Gly Ala Leu His Gly Asp His Gln Arg Phe Ser Thr Phe Ala 225 230 235	782

gga ttc ctg ctc ttt gaa act aag taaatatatg actagaatag ctccactttg 836  
 Gly Phe Leu Leu Phe Glu Thr Lys  
 240 245

gggaagactt gtagctgagc tgatttggtt cgatctgagg aacattaaag ttgagggttt 896  
 tacattgctg tattcaaaaa attattggtt gcaatgttgt tcacgctaca ggtacaccaa 956  
 taatgttgga caattcaggg gctcagaaga atcaaccaca aaatagtctt ctcagatgac 1016  
 ctigactaat atactcagca tctttatcac tctttccttg gcacctaaaa gataattctc 1076  
 ctctgacgca ggttggaat attttttct atcacagaag tcatttgcaa agaattttga 1136  
 ctactctgct tttaatttaa taccagtttt caggaacccc tgaagtttta agttcattat 1196  
 tctttataac atttgagaga atcggatgta gtgatatgac agggctgggg caagaacagg 1256  
 ggcactagct gccttattag ctaatttagt gccctccgtg ttcagcttag cctttgaccc 1316  
 tttccttttg atccacaaaa tacattaaaa ctctgaattc acatacaatg ctattttaaa 1376  
 gtcaatagat tttagctata aagtgttgga ccagtaatgt ggttgtaatt ttgtgtatgt 1436  
 tccccacat cgcccccaac ttcggatgtg gggtcaggag gttgagggtc actattaaca 1496  
 aatgtcataa atatctcata gaggtacagt gccaatagat attcaaagt tgcattgtga 1556  
 ccagagggat tttatatctg aagaacatac actattaata aataccttag agaaagattt 1616  
 tgacctggct ttagataaaa ctgtggcaag aaaaatgtaa tgagcaatat atggaaataa 1676  
 acacaccttt gttaaagata 1696

<210> 2  
 <211> 246  
 <212> PRT  
 <213> Homo sapiens

<400> 2  
 Met Leu Trp Arg Gln Leu Ile Tyr Trp Gln Leu Leu Ala Leu Phe Phe  
 1 5 10 15  
 Leu Pro Phe Cys Leu Cys Gln Asp Glu Tyr Met Glu Ser Pro Gln Thr  
 20 25 30  
 Gly Gly Leu Pro Pro Asp Cys Ser Lys Cys Cys His Gly Asp Tyr Ser  
 35 40 45  
 Phe Arg Gly Tyr Gln Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Ile  
 50 55 60  
 Pro Gly Asn His Gly Asn Asn Gly Asn Asn Gly Ala Thr Gly His Glu  
 65 70 75 80  
 Gly Ala Lys Gly Glu Lys Gly Asp Lys Gly Asp Leu Gly Pro Arg Gly  
 85 90 95  
 Glu Arg Gly Gln His Gly Pro Lys Gly Glu Lys Gly Tyr Pro Gly Ile  
 100 105 110  
 Pro Pro Glu Leu Gln Ile Ala Phe Met Ala Ser Leu Ala Thr His Phe  
 115 120 125

Ser Asn Gln Asn Ser Gly Ile Ile Phe Ser Ser Val Glu Thr Asn Ile  
 130 135 140  
 Gly Asn Phe Phe Asp Val Met Thr Gly Arg Phe Gly Ala Pro Val Ser  
 145 150 155 160  
 Gly Val Tyr Phe Phe Thr Phe Ser Met Met Lys His Glu Asp Val Glu  
 165 170 175  
 Glu Val Tyr Val Tyr Leu Met His Asn Gly Asn Thr Val Phe Ser Met  
 180 185 190  
 Tyr Ser Tyr Glu Met Lys Gly Lys Ser Asp Thr Ser Ser Asn His Ala  
 195 200 205  
 Val Leu Lys Leu Ala Lys Gly Asp Glu Val Trp Leu Arg Met Gly Asn  
 210 215 220  
 Gly Ala Leu His Gly Asp His Gln Arg Phe Ser Thr Phe Ala Gly Phe  
 225 230 235 240  
 Leu Leu Phe Glu Thr Lys  
 245

&lt;210&gt; 3

&lt;211&gt; 244

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

Met Leu Leu Leu Gly Ala Val Leu Leu Leu Leu Ala Leu Pro Gly His  
 1 5 10 15  
 Asp Gln Glu Thr Thr Thr Gln Gly Pro Gly Val Leu Leu Pro Leu Pro  
 20 25 30  
 Lys Gly Ala Cys Thr Gly Trp Met Ala Gly Ile Pro Gly His Pro Gly  
 35 40 45  
 His Asn Gly Ala Pro Gly Arg Asp Gly Arg Asp Gly Thr Pro Gly Glu  
 50 55 60  
 Lys Gly Glu Lys Gly Asp Pro Gly Leu Ile Gly Pro Lys Gly Asp Ile  
 65 70 75 80  
 Gly Glu Thr Gly Val Pro Gly Ala Glu Gly Pro Arg Gly Phe Pro Gly  
 85 90 95  
 Ile Gln Gly Arg Lys Gly Glu Pro Gly Glu Gly Ala Tyr Val Tyr Arg  
 100 105 110  
 Ser Ala Phe Ser Val Gly Leu Glu Thr Tyr Val Thr Ile Pro Asn Met  
 115 120 125  
 Pro Ile Arg Phe Thr Lys Ile Phe Tyr Asn Gln Gln Asn His Tyr Asp  
 130 135 140  
 Gly Ser Thr Gly Lys Phe His Cys Asn Ile Pro Gly Leu Tyr Tyr Phe  
 145 150 155 160

[illegible]

```
<210> 4
<211> 245
<212> PRT
<213> Homo sapiens
```

<400> 4															
Met	Asp	Val	Gly	Pro	Ser	Ser	Leu	Pro	His	Leu	Gly	Leu	Lys	Leu	Leu
1				5					10					15	
Leu	Leu	Leu	Leu	Leu	Leu	Ala	Leu	Arg	Gly	Gln	Ala	Asn	Thr	Gly	Cys
			20					25					30		
Tyr	Gly	Ile	Pro	Gly	Met	Pro	Gly	Leu	Pro	Gly	Ala	Pro	Gly	Lys	Asp
		35					40					45			
Gly	Tyr	Asp	Gly	Leu	Pro	Gly	Pro	Lys	Gly	Glu	Pro	Gly	Ile	Pro	Ala
	50					55					60				
Ile	Pro	Gly	Ile	Arg	Gly	Pro	Lys	Gly	Gln	Lys	Gly	Glu	Pro	Gly	Leu
65					70					75					80
Pro	Gly	His	Pro	Gly	Lys	Asn	Gly	Pro	Met	Gly	Pro	Pro	Gly	Met	Pro
				85					90					95	
Gly	Val	Pro	Gly	Pro	Met	Gly	Ile	Pro	Gly	Glu	Pro	Gly	Glu	Glu	Gly
			100					105					110		
Arg	Tyr	Lys	Gln	Lys	Phe	Gln	Ser	Val	Phe	Thr	Val	Thr	Arg	Gln	Thr
		115					120					125			
His	Gln	Pro	Pro	Ala	Pro	Asn	Ser	Leu	Ile	Arg	Phe	Asn	Ala	Val	Leu
	130					135						140			
Thr	Asn	Pro	Gln	Gly	Asp	Tyr	Asp	Thr	Ser	Thr	Gly	Lys	Phe	Thr	Cys
145					150					155					160
Lys	Val	Pro	Gly	Leu	Tyr	Tyr	Phe	Val	Tyr	His	Ala	Ser	His	Thr	Ala
				165					170					175	
Asn	Leu	Cys	Val	Leu	Leu	Tyr	Arg	Ser	Gly	Val	Lys	Val	Val	Thr	Phe
			180					185					190		

Cys	Gly	His	Thr	Ser	Lys	Thr	Asn	Gln	Val	Asn	Ser	Gly	Gly	Val	Leu
		195					200					205			
Leu	Arg	Leu	Gln	Val	Gly	Glu	Glu	Val	Trp	Leu	Ala	Val	Asn	Asp	Tyr
	210					215					220				
Tyr	Asp	Met	Val	Gly	Ile	Gln	Gly	Ser	Asp	Ser	Val	Phe	Ser	Gly	Phe
225					230					235					240
Leu	Leu	Phe	Pro	Asp											
				245											

<210> 5

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Aromatic motif

<221> VARIANT

<222> (2)...(6)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (7)...(7)

<223> Xaa is asparagine or aspartic acid

<221> VARIANT

<222> (8)...(11)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (12)...(12)

<223> Xaa is phenylalanine, tyrosine, tryptophan or leucine

<221> VARIANT

<222> (13)...(18)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (20)...(24)

<223> Each Xaa is independently any amino acid residue.

7

<221> VARIANT  
 <222> (26)...(26)  
 <223> Xaa is any amino acid residue

<221> VARIANT  
 <222> (28)...(28)  
 <223> Xaa is any amino acid residue

<221> VARIANT  
 <222> (30)...(30)  
 <223> Xaa is any amino acid residue

<221> VARIANT  
 <222> (31)...(31)  
 <223> Xaa is phenylalanine or tyrosine

<400> 5  
 Phe Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 1 5 10 15  
 Xaa Xaa Phe Xaa Xaa Xaa Xaa Xaa Gly Xaa Tyr Xaa Phe Xaa Xaa  
 20 25 30

<210> 6  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Degenerate nucleotide primer

<221> variation  
 <222> (1)...(18)  
 <223> Each N is independently any nucleotide

<400> 6  
 ggngansarg tntggyt

17

<210> 7  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
<223> Degenerate nucleotide primer

<221> variation  
<222> (1)...(17)  
<223> Each N is independently any nucleotide

<400> 7  
sngnnntnta ytwyttyr 18

<210> 8  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Degenerate nucleotide primer

<221> variation  
<222> (1)...(17)  
<223> Each N is independently any nucleotide

<400> 8  
ttydsnggnt tyytnht 17

<210> 9  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Degenerate nucleotide primer

<221> variation  
<222> (1)...(18)  
<223> Each N is independently any nucleotide

<400> 9  
ytwyrayrbn wbnwsngg 18

<210> 10  
<211> 738  
<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate nucleotide sequence encoding the polypeptide of SEQ ID NO:2

<221> variation

<222> (1)...(738)

<223> Each N is independently any nucleotide

<400> 10

atgytntggm gncarytnat htaytggcar ytnytngcny tnttyttyyt nccnttytgy	60
ytntgycarg aygartayat ggarwsnccn caracngng gnytnccncc ngaytgywsn	120
aartgytgyc ayggngayta ywsnttymgn ggntaycarg gncncncngg nccncncngn	180
ccncncngna thccnggnaa ycayggnay aayggnaaya ayggngcnac nggncaygar	240
ggngcnaarg gngaraarg ngayaarggn gayytngnc cnmgngnga rmngngncar	300
cayggncna arggngaraa rggntayccn ggnathccnc cngarytnca rathgcntty	360
atggcnwsny tngcnacna yttywsnaay caraaywsng gnathathtt ywsnwsngtn	420
garacnaaya thggnaaytt ytygaygt n atgacnggm gnttyggngc nccngtnwsn	480
ggngtntayt tyttyacntt ywsnatgat aarcaygarg aygtngarga rgtntaygt n	540
taytntatgc ayaayggnaa yacngtntty wsnatgtayw sntaygarat gaarggnaar	600
wsngayacnw snwsnaayca ygcngtnytn aarytngcna arggngayga rgtntggytn	660
mgnatgggna ayggngcnyt ncayggngay caycarmgnt tywsnacntt ygcnggntty	720
ytnytnttyg aracnaar	738

<210> 11

<211> 1117

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (111)...(848)

<400> 11

gctgcagctc tcattctcaa acctggcatt tgctgaggc gaccacggta cctccagccc	60
ctgtcaagct tccctgcgag actcttgctg atttgccgat ttgccgagcc atg ctc	116
	Met Leu
	1
ggg agg cag cgc atc tgg tgg cac ctg ctg cct ttg ctt ttc ctc cca	164
Gly Arg Gln Arg Ile Trp Trp His Leu Leu Pro Leu Leu Phe Leu Pro	
5 10 15	

ttt tgc ctg tgt caa gat gaa tac atg gag tct cca caa gct gga gga Phe Cys Leu Cys Gln Asp Glu Tyr Met Glu Ser Pro Gln Ala Gly Gly 20 25 30	212
ctg ccc cca gac tgc agc aag tgt tgc cat gga gat tat gga ttt cgt Leu Pro Pro Asp Cys Ser Lys Cys Cys His Gly Asp Tyr Gly Phe Arg 35 40 45 50	260
ggt tac caa ggg ccc cct gga cct cca ggt cct cct ggc att cca gga Gly Tyr Gln Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Ile Pro Gly 55 60 65	308
aac cat gga aac aat ggg aac aat gga gct act ggc cat gaa ggg gcc Asn His Gly Asn Asn Gly Asn Asn Gly Ala Thr Gly His Glu Gly Ala 70 75 80	356
aaa ggt gag aaa gga gac aaa ggc gac cta ggc cct cga gga gaa cgg Lys Gly Glu Lys Gly Asp Lys Gly Asp Leu Gly Pro Arg Gly Glu Arg 85 90 95	404
ggg cag cat ggc ccc aaa gga gag aaa ggc tac cca ggg gtg cca cca Gly Gln His Gly Pro Lys Gly Glu Lys Gly Tyr Pro Gly Val Pro Pro 100 105 110	452
gaa ctg cag att gca ttc atg gct tct cta gca act cac ttc agc aat Glu Leu Gln Ile Ala Phe Met Ala Ser Leu Ala Thr His Phe Ser Asn 115 120 125 130	500
cag aac agt ggc att atc ttc agc agt gtt gag acc aac att gga aac Gln Asn Ser Gly Ile Ile Phe Ser Ser Val Glu Thr Asn Ile Gly Asn 135 140 145	548
ttc ttc gat gtc atg act ggg aga ttt ggg gcc ccc gta tca ggt gtg Phe Phe Asp Val Met Thr Gly Arg Phe Gly Ala Pro Val Ser Gly Val 150 155 160	596
tat ttc ttc acc ttc agc atg atg aag cat gag gac gta gag gaa gtg Tyr Phe Phe Thr Phe Ser Met Met Lys His Glu Asp Val Glu Glu Val 165 170 175	644
tat gtg tac ctt atg cac aac ggc aac aca gtc ttc agc atg tac agc Tyr Val Tyr Leu Met His Asn Gly Asn Thr Val Phe Ser Met Tyr Ser	692

180	185	190	
tat gaa aca aag gga aaa tca gat aca tcc agc aac cat gca gtg ctg			740
Tyr Glu Thr Lys Gly Lys Ser Asp Thr Ser Ser Asn His Ala Val Leu			
195	200	205	210
aag ttg gcc aaa gga gat gaa gtc tgg cta aga atg ggc aac ggt gcc			788
Lys Leu Ala Lys Gly Asp Glu Val Trp Leu Arg Met Gly Asn Gly Ala			
	215	220	225
ctc cac ggg gac cac cag cgc ttc tcc acc ttc gca ggc ttt ctg ctc			836
Leu His Gly Asp His Gln Arg Phe Ser Thr Phe Ala Gly Phe Leu Leu			
	230	235	240
ttt gaa act aag tgacaaggaa gacaggatat ctccactttg ggggcaattt			888
Phe Glu Thr Lys			
245			
atagctgagc tagggttggt aggatatgaa ggatgttgaa gtcggggggt ctttatggag			948
catttaagtg ttgcattggt cacactgcta ctcatcttaa tggcatacca ataatgttgg			1008
atgcttcagg ggctcactgc tactcattct aatggcatac caataatgtt ggatgcttca			1068
ggggctcact gctactcatt ctaatggcaa taccaataat gttggatgc			1117

&lt;210&gt; 12

&lt;211&gt; 246

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 12

Met Leu Gly Arg Gln Arg Ile Trp Trp His Leu Leu Pro Leu Leu Phe	
1 5 10 15	
Leu Pro Phe Cys Leu Cys Gln Asp Glu Tyr Met Glu Ser Pro Gln Ala	
20 25 30	
Gly Gly Leu Pro Pro Asp Cys Ser Lys Cys Cys His Gly Asp Tyr Gly	
35 40 45	
Phe Arg Gly Tyr Gln Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly Ile	
50 55 60	
Pro Gly Asn His Gly Asn Asn Gly Asn Asn Gly Ala Thr Gly His Glu	
65 70 75 80	
Gly Ala Lys Gly Glu Lys Gly Asp Lys Gly Asp Leu Gly Pro Arg Gly	
85 90 95	
Glu Arg Gly Gln His Gly Pro Lys Gly Glu Lys Gly Tyr Pro Gly Val	
100 105 110	

12

Pro Pro Glu Leu Gln Ile Ala Phe Met Ala Ser Leu Ala Thr His Phe  
           115                          120                          125  
 Ser Asn Gln Asn Ser Gly Ile Ile Phe Ser Ser Val Glu Thr Asn Ile  
           130                          135                          140  
 Gly Asn Phe Phe Asp Val Met Thr Gly Arg Phe Gly Ala Pro Val Ser  
 145                          150                          155                          160  
 Gly Val Tyr Phe Phe Thr Phe Ser Met Met Lys His Glu Asp Val Glu  
                           165                          170                          175  
 Glu Val Tyr Val Tyr Leu Met His Asn Gly Asn Thr Val Phe Ser Met  
                           180                          185                          190  
 Tyr Ser Tyr Glu Thr Lys Gly Lys Ser Asp Thr Ser Ser Asn His Ala  
           195                          200                          205  
 Val Leu Lys Leu Ala Lys Gly Asp Glu Val Trp Leu Arg Met Gly Asn  
           210                          215                          220  
 Gly Ala Leu His Gly Asp His Gln Arg Phe Ser Thr Phe Ala Gly Phe  
 225                          230                          235                          240  
 Leu Leu Phe Glu Thr Lys  
                           245

&lt;210&gt; 13

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Oligonucleotide ZC21913

&lt;400&gt; 13

tgaccagagg gattttat

18

&lt;210&gt; 14

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Oligonucleotide ZC21914

&lt;400&gt; 14

ttgccacagt tttatcta

18

&lt;210&gt; 15

&lt;211&gt; 25

13

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide ZC23377

<400> 15  
ctctgttggg atccatgctt tgga

24

<210> 16  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> ZC23378

<400> 16  
gtcatctaga tacttagttt ca

22

<210> 17  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Glu-Glu tag

<400> 17  
Glu Glu Tyr Met Pro Met Glu  
1 5

<210> 18  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> ZC447

<400> 18  
taacaatttc acacagg

17

14

<210> 19  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide ZC976

<400> 19  
cgttgtaaaa cgacggcc

18

<210> 20  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> purification peptide

<400> 20  
Glu Tyr Met Pro Val Asp  
1 5

# INTERNATIONAL SEARCH REPORT

Int lional Application No

PCT/US 00/10454

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C12N15/62 C07K16/18 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, STRAND, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 06551 A (LACROIX BRUNO ;DUCLERT AYMERIC (FR); GENSET (FR); DUMAS MILNE EDWA) 11 February 1999 (1999-02-11)	12,31
A	Sequences ID no.307 and no.149	1-11,13, 21-30, 32-48
X	NCI-CGAP: "National cancer institute, cancer genome anatomy project (CGAP)" EMBL DATABASE ENTRY HS1318772, ACCESSION NUMBER AA527298, 22 July 1997 (1997-07-22), XP002144457 "100% identity in 594 bp overlap with sequence ID no.1 from 168" & UNPUBLISHED,	31

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 August 2000

Date of mailing of the international search report

23/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Le Cornec, N

## INTERNATIONAL SEARCH REPORT

In tional Application No  
PCT/US 00/10454

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	M.D. ADAMS ET AL: "Initial assessment of human gene diversity and expression patterns based upon 83 million basepairs of cDNA sequence " EMBL DATABASE ENTRY HSZZ39660, ACCESSION NUMBER AA334609, 21 April 1997 (1997-04-21), XP002144458 "EST38850: 100% identity in 355 bp overlap with sequence ID no.1" abstract & NATURE, vol. 377, 1995, pages 3-174, ---	31
A	WO 96 39429 A (WHITEHEAD BIOMEDICAL INST ;SCHERER PHILIPP E (US); LODISH HARVEY F) 12 December 1996 (1996-12-12) cited in the application claims; figures 1,2 ---	1-48
A	WO 99 07736 A (BOUGUELERET LYDIE ;INST NAT SANTE RECH MED (FR); YEN POTIN FRANCES) 18 February 1999 (1999-02-18) Sequence ID no.1 ---	1,4
A	MAEDA K ET AL: "cDNA CLONING AND EXPRESSION OF A NOVEL ADIPOSE SPECIFIC COLLAGEN- LIKE FACTOR, APM1 (ADIPOSE MOST ABUNDANT GENE TRANSCRIPT)" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS,US,ACADEMIC PRESS INC. ORLANDO, FL, vol. 221, 1 April 1996 (1996-04-01), pages 286-289, XP000612064 ISSN: 0006-291X cited in the application the whole document ---	1-48
A	SELLAR GC ET AL: "Characterization and organization of the genes encoding the A-, B- and C-chains of human complement subcomponent Clq" BIOCHEMICAL JOURNAL,GB,PORTLAND PRESS, LONDON, vol. 274, 1 March 1991 (1991-03-01), pages 481-490, XP002087332 ISSN: 0264-6021 cited in the application the whole document -----	1-48

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/10454

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9906551	A	11-02-1999	AU	8555498 A	22-02-1999
			EP	1000149 A	17-05-2000
WO 9639429	A	12-12-1996	US	5869330 A	09-02-1999
WO 9907736	A	18-02-1999	FR	2767136 A	12-02-1999
			FR	2767135 A	12-02-1999
			AU	8556398 A	01-03-1999
			AU	8556498 A	01-03-1999
			EP	1003783 A	31-05-2000
			EP	0964870 A	22-12-1999
			WO	9907737 A	18-02-1999

**THIS PAGE BLANK (USPTO)**